



SUMMARY OF PRODUCT CHARACTERISTIC

1. Name of the medicinal product

Bima-opt

2. Qualitative and quantitative declaration

One ml of solution contains 0.1 mg bimatoprost.

Excipient with known effect:

One ml of solution contains 0.2 mg benzalkonium chloride.

For the full list of excipients, see section 6.1.

3. Pharmaceutical form

Eye drops, solution.

Eye drop multiple dose.

Clear colorless sterile solution

4. Clinical particulars

4.1 Therapeutic indications

Reduction of elevated intraocular pressure in chronic open-angle glaucoma and ocular hypertension in adults (as monotherapy or as adjunctive therapy to beta-blockers)

4.2 Posology and method of administration

Posology

The recommended dose is one drop in the affected eye(s) once daily, administered in the evening. The dose should not exceed once daily, as more frequent administration may lessen the intraocular pressure lowering effect.

Pediatric population

The safety and efficacy of bimatoprost in children aged 0 to 18 years has not yet been established.

Patients with hepatic and renal impairment

Bimatoprost has not been studied in patients with renal or moderate to severe hepatic impairment and should therefore be used with caution in such patients. In patients with a history of mild liver disease or abnormal alanine aminotransferase (ALT), aspartate aminotransferase (AST) and/or bilirubin at baseline, bimatoprost 0.3 mg/ml eye drops solution had no adverse effect on liver function over 24 months.

Method of administration

If more than one topical ophthalmic medicinal product is being used, each one should be administered at least 5 minutes apart.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Bima-opt 0.1 mg/ml is contraindicated in patients who have had a suspected previous adverse reaction to benzalkonium chloride that has led to discontinuation.

4.4 Special warnings and precautions for use

Ocular

Before treatment is initiated, patients should be informed of the possibility of prostaglandin analogue periorbitopathy (PAP) and increased iris pigmentation, since these have been observed during treatment with bimatoprost. Some of these changes may be permanent, and may lead to impaired field of vision and differences in appearance between the eyes when only one eye is treated (see section 4.8).

Cystoid macular edema has been uncommonly reported ($\geq 1/1,000$ to $< 1/100$) following treatment with bimatoprost 0.3 mg/ml eye drops, solution. Therefore, bimatoprost 0.3 mg should be used with caution in patients with known risk factors for macular edema (e.g. aphakic patients, pseudophakic patients with a torn posterior lens capsule).

There have been rare spontaneous reports of reactivation of previous corneal infiltrates or ocular infections with bimatoprost 0.3 mg/ml eye drops, solution. Bimatoprost should be used with caution in patients with a prior history of significant ocular viral infections (e.g. herpes simplex) or uveitis/iritis.

Bimatoprost has not been studied in patients with inflammatory ocular conditions, neovascular, inflammatory, angle-closure glaucoma, congenital glaucoma or narrow-angle glaucoma.

Skin

There is a potential for hair growth to occur in areas where bimatoprost solution comes repeatedly in contact with the skin surface. Thus, it is important to apply bimatoprost as instructed and avoid it running onto the cheek or other skin areas.

Respiratory

Bimatoprost has not been studied in patients with compromised respiratory function. While there is limited information available on patients with a history of asthma or COPD, there have been reports of exacerbation of asthma, dyspnea and COPD, as well as reports of asthma, in post marketing experience. The frequency of these symptoms is not known. Patients with COPD, asthma or compromised respiratory function due to other conditions should be treated with caution.

Cardiovascular

Bimatoprost has not been studied in patients with heart block more severe than first degree or uncontrolled congestive heart failure. There have been a limited number of spontaneous reports of bradycardia or hypotension



with bimatoprost 0.3 mg/ml eye drops solution. It should be used with caution in patients predisposed to low heart rate or low blood pressure.

Other information

In studies of bimatoprost 0.3 mg/ml in patients with glaucoma or ocular hypertension, it has been shown that the more frequent exposure of the eye to more than one dose of bimatoprost daily may decrease the IOP-lowering effect (see section 4.5). Patients using bimatoprost with other prostaglandin analogues should be monitored for changes to their intraocular pressure.

Bima-opt 0.1 mg/ml contains the preservative benzalkonium chloride (200 ppm), which may be absorbed by soft contact lenses. Eye irritation and discoloration of the soft contact lenses may also occur because of the presence of benzalkonium chloride. Contact lenses should be removed prior to instillation and may be reinserted 15 minutes following administration.

Benzalkonium chloride, which is commonly used as a preservative in ophthalmic products, has been reported to cause punctate keratopathy and/or toxic ulcerative keratopathy. It should be used with caution in dry eye patients, in patients where the cornea may be compromised and in patients taking multiple BAK-containing eye drops. In addition, monitoring is required with prolonged use in such patients.

There have been reports of bacterial keratitis associated with the use of multiple dose containers of topical ophthalmic products. These containers had been inadvertently contaminated by patients who, in most cases, had a concurrent ocular disease. Patients with a disruption of the ocular epithelial surface are at greater risk of developing bacterial keratitis.

Patients should be instructed to avoid allowing the tip of the dispensing container to contact the eye or surrounding structures, to avoid eye injury and contamination of the solution.

4.5 Interaction with other medicinal products and other forms of interaction

No interaction studies have been performed.

No interactions are anticipated in humans, since systemic concentrations of bimatoprost are extremely low (less than 0.2 ng/ml) following ocular dosing with bimatoprost 0.3 mg/ml eye drops, solution. Bimatoprost is biotransformed by any of multiple enzymes and pathways, and no effects on hepatic drug metabolizing enzymes were observed in preclinical studies.

In clinical studies, bimatoprost 0.3 mg/ml, eye drops solution was used concomitantly with a number of different ophthalmic beta-blocking agents without evidence of interactions.

Concomitant use of bimatoprost and antiglaucomatous agents other than topical beta-blockers has not been evaluated during adjunctive glaucoma therapy.

Only a potential for the IOP-lowering effect of prostaglandin analogues was reduced in patients with glaucoma or ocular hypertension when used with other prostaglandin analogues (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no adequate data from the use of bimatoprost in pregnant women. Animal studies have shown reproductive toxicity at high maternotoxic doses (see section 5.3).

It should not be used during pregnancy unless clearly necessary.

Breast-feeding

It is unknown whether bimatoprost is excreted in human breast milk. Animal studies have shown excretion of bimatoprost in breast milk. A decision must be made whether to discontinue breast-feeding or to discontinue from bima-opt therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Fertility - There are no data on the effects of bimatoprost on human fertility.

4.7 Effects on ability to drive and use machines

Bima-opt has negligible influence on the ability to drive and use machines. As with any ocular treatment, if transient blurred vision occurs at instillation, the patient should wait until the vision clears before driving or using machines.

4.8 Undesirable effects

In a 12-month Phase III clinical study approximately 38 % of patients treated with bimatoprost 0.1 mg/ml eye drops, solution experienced adverse reactions. The most frequently reported adverse reaction was conjunctival hyperaemia (mostly trace to mild and of a non-inflammatory nature) occurring in 29 % of patients. Approximately 4 % of patients discontinued due to any adverse event in the 12-month study.

The following adverse reactions were reported during clinical trials with bimatoprost 0.1 mg/ml eye drops solution or in the post-marketing period. Most were ocular, mild and none was serious.

Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); not known (cannot be estimated from available data) adverse reactions are presented according to System Organ Class in table 1 in order of decreased seriousness within each frequency grouping.

Table 1.

System Organ class	Frequency	Adverse reaction
<i>Nervous system disorders</i>	uncommon	Headache
	not known	Dizziness
<i>Respiratory, thoracic and mediastinal disorders</i>	not known	Asthma, asthma exacerbation, COPD exacerbation and dyspnea

System Organ class	Frequency	Adverse reaction
<i>Eye disorders</i>	very common	Conjunctiva hyperemia, prostaglandin analogue periorbitopathy
	common	Punctate keratitis, eye irritation, eye pruritus, growth of eyelashes, eye pain, erythema of eyelid, eyelid pruritus
	uncommon	Asthenopia, blurred vision, conjunctiva disorder, conjunctiva edema, iris hyperpigmentation, madarosis, eyelid edema
	not known	Blepharal pigmentation, macular edema, dry eye, eye discharge, eye edema, foreign body sensation in eyes, lacrimation increased, ocular discomfort, photophobia
<i>Gastrointestinal disorders</i>	uncommon	Nausea
<i>Skin and subcutaneous tissue disorders</i>	common	Skin hyperpigmentation, hypertrichosis
	uncommon	Dry skin, eyelid margin crusting, pruritus
	not known	Skin discoloration (periocular)
<i>General disorders and administration site conditions</i>	common	Instillation site irritation
<i>Immune system disorders</i>	not known	Hypersensitivity reaction including signs and symptoms of eye allergy and allergic dermatitis
<i>Vascular disorders</i>	not known	Hypertension

Description of selected adverse reactions

Prostaglandin analogue periorbitopathy (PAP)

Prostaglandin analogues including bimatoprost can induce periorbital lipodystrophic changes which can lead to deepening of the eyelid sulcus, ptosis, enophthalmos, eyelid retraction, involution of dermatochalasis and inferior scleral show. Changes are typically mild, can occur as early as one month after initiation of treatment with bima-opt, and may cause impaired field of vision even in the absence of patient recognition. PAP is also associated with periocular skin hyperpigmentation or discoloration and hypertrichosis. All changes have been noted to be partially or fully reversible upon discontinuation or switch to alternative treatments.

Iris hyperpigmentation

Increased iris pigmentation is likely to be permanent. The pigmentation change is due to increased melanin content in the melanocytes rather than to an increase in the number of melanocytes. The long-term effects of increased iris pigmentation are not known. Iris color changes seen with ophthalmic administration of bimatoprost may not be noticeable for several months to years. Typically, the brown pigmentation around the pupil spreads concentrically towards the periphery of the iris and the entire iris or parts become more brownish. Neither naevi nor freckles of the iris appear to be affected by the treatment. At 12 months, the incidence of iris hyperpigmentation with bimatoprost 0.1 mg/ml eye drops, solution was 0.5%. At 12 months, the incidence with bimatoprost 0.3 mg/ml eye drops, solution was 1.5% (see section 4.8 Table 2) and did not increase following 3 years treatment.

In clinical studies, over 1800 patients have been treated with bimatoprost 0.3 mg/ml eye drops, solution. On combining the data from phase III monotherapy and adjunctive bimatoprost 0.3 mg/ml eye drops, solution usage, the most frequently reported adverse events were:

- Growth of eyelashes in up to 45% in the first year with the incidence of new reports decreasing to 7% at 2 years and 2% at 3 years
- Conjunctival hyperemia (mostly trace to mild and thought to be of a non-inflammatory nature) in up to 44% in the first year with the incidence of new reports decreasing to 13% at 2 years and 12% at 3 years
- Ocular pruritus in up to 14% of patients in the first year with the incidence of new reports decreasing to 3% at 2 years and 0% at 3 years. Less than 9% of patients discontinued due to any adverse event in the first year with the incidence of additional patient discontinuations being 3% at both 2 and 3 years.

Additional adverse reactions reported with bimatoprost 0.3 mg/ml are presented in Table 2. The table also includes those adverse reactions which occurred with both formulations but at a different frequency. Most were ocular, mild to moderate, and none was serious: With each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Table 2.

System Organ class	Frequency	Adverse reaction
<i>Nervous system disorders</i>	common	Headache
	uncommon	Dizziness
<i>Eye disorders</i>	very common	Ocular pruritus, growth of eyelashes
	common	Corneal erosion, ocular burning, allergic conjunctivitis, blepharitis, worsening of visual acuity, asthenopia, conjunctival edema, foreign body sensation, ocular

		dryness, eye pain, photophobia, tearing, eye discharge, visual disturbance/blurred vision, increased iris pigmentation, eyelash darkening
	uncommon	Retinal hemorrhage, uveitis, cystoid macular edema, iritis, blepharospasm, eyelid retraction, periorbital erythema
<i>Skin and subcutaneous tissue disorders</i>	uncommon	Hirsutism
<i>Vascular disorders</i>	common	Hypertension
<i>General disorders and administration site conditions</i>	uncommon	Asthenia
<i>Investigations</i>	common	Liver function test abnormal

Adverse reactions reported in phosphate containing eye drops: Cases of corneal calcification have been reported very rarely in association with the use of phosphate containing eye drops in some patients with significantly damaged corneas.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Health Product Pharmacovigilance Center at <http://thaihpvc.fda.moph.go.th>

4.9 Overdose

No case of overdose has been reported, and is unlikely to occur after ocular administration.

If overdose occurs, treatment should be symptomatic and supportive. If bima-opt is accidentally ingested, the following information may be useful: in two-week oral rat and mouse studies, doses up to 100 mg/kg/day did not produce any toxicity. This dose expressed as mg/m² is at least 210 times higher than the accidental dose of one bottle of bimatoprost 0.1 mg/ml eye drops, solution in a 10 kg child.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Ophthalmologicals, prostaglandin analogues, ATC code: S01EE03



Mechanism of action

The mechanism of action by which bimatoprost reduces intraocular pressure in humans is by increasing aqueous humor outflow through the trabecular meshwork and enhancing uveoscleral outflow. Reduction of the intraocular pressure starts approximately 4 hours after the first administration and maximum effect is reached within approximately 8 to 12 hours. The duration of effect is maintained for at least 24 hours.

Bimatoprost is a potent ocular hypotensive agent. It is a synthetic prostamide, structurally related to prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$), that does not act through any known prostaglandin receptors. Bimatoprost selectively mimics the effects of newly discovered biosynthesized substances called prostamides. The prostamide receptor, however, has not yet been structurally identified.

During a 12-month pivotal study in adults with bimatoprost 0.1 mg/ml eye drops, the mean diurnal IOP values measured at any visit over the 12-month study period differed by no more than 1.1 mmHg throughout the day and were never greater than 17.7 mmHg.

Bimatoprost 0.1 mg/ml eye drops contains BAK in a concentration of 200 ppm.

Limited experience is available in patients with open-angle glaucoma with pseudoexfoliative and pigmentary glaucoma, and chronic angle-closure glaucoma with patent iridotomy.

No clinically relevant effects on heart rate and blood pressure have been observed in clinical trials.

Pediatric population

The safety and efficacy of bimatoprost in children aged 0 to less than 18 years has not been established.

5.2 Pharmacokinetic properties

Absorption

Bimatoprost penetrates the human cornea and sclera well *in vitro*. After ocular administration in adults, the systemic exposure of bimatoprost is very low with no accumulation over time. After once daily ocular administration of one drop of 0.3 mg/ml bimatoprost to both eyes for two weeks, blood concentrations peaked within 10 minutes after dosing and declined to below the lower limit of detection (0.025 ng/ml) within 1.5 hours after dosing. Onset generally occurs within 4 hours after application to the eyes and peaks within 8-12 hours. Mean C_{max} and $AUC_{0-24hrs}$ values were similar on days 7 and 14 at approximately 0.08 ng/ml and 0.09 ng•hr/ml respectively, indicating that a steady bimatoprost concentration was reached during the first week of ocular dosing.

Distribution

Bimatoprost is moderately distributed into body tissues and the systemic volume of distribution in humans at steady-state was 0.67 l/kg. In human blood, bimatoprost resides mainly in the plasma. The plasma protein binding of bimatoprost is approximately 88 %.

Metabolism

Bimatoprost is the major circulating species in the blood once it reaches the systemic circulation following ocular dosing. It then undergoes oxidation, N-deethylation and glucuronidation to form various metabolites.

Elimination

Bimatoprost is eliminated primarily by renal excretion, up to 67% of an intravenous dose administered to healthy adult volunteers was excreted in the urine, 25% of the dose was excreted via the feces. The elimination half-life, determined after intravenous administration, was approximately 45 minutes; the total blood clearance was 1.5 l/hr/kg.

Characteristics in elderly patients

After twice daily dosing with bimatoprost 0.3 mg/ml eye drops, solution, the mean AUC_{0-24hr} value of 0.0634 ng•hr/ml bimatoprost in the elderly (subjects 65 years or older) were significantly higher than 0.0218 ng•hr/ml in young healthy adults. However, this finding is not clinically relevant as systemic exposure for both elderly and young subjects remained very low from ocular dosing. There was no accumulation of bimatoprost in the blood over time and the safety profile was similar in elderly and young patients.

5.3 Preclinical safety data

Effects in non-clinical studies were observed only at exposures considered sufficiently in excess of the maximum human exposure indicating little relevance to clinical use.

Monkeys administered ocular bimatoprost concentrations of ≥ 0.3 mg/ml daily for 1 year had an increase in iris pigmentation and reversible dose-related periocular effects characterized by a prominent upper and/or lower sulcus and widening of the palpebral fissure. The increased iris pigmentation appears to be caused by increased stimulation of melanin production in melanocytes and not by an increase in melanocyte number. No functional or microscopic changes related to the periocular effects have been observed, and the mechanism of action for the periocular changes is unknown.

Bimatoprost was not mutagenic or carcinogenic in a series of *in vitro* and *in vivo* studies.

Bimatoprost did not impair fertility in rats up to doses of 0.6 mg/kg/day (at least 103-times the intended human exposure). In embryo/fetal developmental studies abortion, but no developmental effects were seen in mice and rats at doses that were at least 860-times or 1700-times higher than the dose in humans, respectively. These doses resulted in systemic exposures of at least 33- or 97-times higher, respectively, than the intended human exposure. In rat peri/postnatal studies, maternal toxicity caused reduced gestation time, fetal death, and decreased pup body weights at ≥ 0.3 mg/kg/day (at least 41-times the intended human exposure). Neurobehavioral functions of offspring were not affected.



PHARMA INNOVA CO., LTD.

1/38 Moo 4, Liebkhlong 7 Road, Buengkamproi, Lam Luk Ka, Pathumthani 12150, Thailand

Tel. (662) 971-5335 Fax (662) 971-5470 E-mail: info@pharmainnova.co.th

6. Pharmaceutical particulars

6.1 List of excipients

- Sodium chloride
- Dibasic sodium phosphate
- Citric acid
- Benzalkonium chloride
- Water for injection q.s. to
- Sodium hydroxide (10%w/v) and/or
- Hydrochloric acid (10%w/v) q.s. to pH 6.8-7.8

6.2 Incompatibilities

Not applicable

6.3 Shelf life

2 years

After first opening: 30 days

6.4 Special precautions for storage

Store below 30°C

6.5 Nature and contents of container

Plastic bottle (LDPE) of 3 mL into paper box of 1, 10, 20, 50, and 100 bottles

6.6 Special precautions for disposal and other handling

No special requirements for disposal.

7. Marketing authorisation holder

PHARMA INNOVA CO., LTD.

1/38 Moo 4, Liebkhlong 7 Road, Buengkamproi,

Lam Luk Ka, Pathumthani 12150, Thailand

Tel. 0-2532-7181 Fax 0-2532-7019

8. Marketing authorisation number(s)

xx xxx/xx

9. Date of first authorization/ renewal of the authorization

DD/MM/YYYY

10. Date of revision of the text

7 October 2022