

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

NovoEight 250 IU powder and solvent for solution for injection
NovoEight 500 IU powder and solvent for solution for injection

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

NovoEight 250 IU powder and solvent for solution for injection.

Each powder vial contains nominally 250 IU human coagulation factor VIII (rDNA), turoctocog alfa.

After reconstitution NovoEight contains approximately 62.5 IU/ml of human coagulation factor VIII (rDNA), turoctocog alfa.

NovoEight 500 IU powder and solvent for solution for injection.

Each powder vial contains nominally 500 IU human coagulation factor VIII (rDNA), turoctocog alfa.

After reconstitution NovoEight contains approximately 125 IU/ml of human coagulation factor VIII (rDNA), turoctocog alfa.

The potency (IU) is determined using the European Pharmacopoeia chromogenic assay. The specific activity of NovoEight is approximately 8,300 IU/mg protein.

Turoctocog alfa (human coagulation factor VIII (rDNA)) is a purified protein that has 1,445 amino acids with an approximate molecular mass of 166 kDA. It is produced by recombinant DNA technology in Chinese hamster ovary (CHO) cells, and prepared without the addition of any human or animal derived protein in the cell culture process, purification or final formulation.

Turoctocog alfa is a B-domain truncated recombinant human coagulation factor VIII (B-domain consists of 21 amino acids of the wild type B-domain) without any other modifications in the amino acid sequence.

Excipient with known effect

0.31 mmol sodium (7 mg) per ml of reconstituted solution.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder and solvent for solution for injection.

White or slightly yellow powder or friable mass.

Clear and colourless solution for injection.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Treatment and prophylaxis of bleeding in patients with haemophilia A (congenital factor VIII deficiency).

4.2 Posology and method of administration

Treatment should be initiated under the supervision of a doctor experienced in the treatment of haemophilia.

Posology

The dosage and duration of the substitution therapy depend on the severity of the factor VIII deficiency, on the location and extent of the bleeding and the patient's clinical condition.

The number of units of factor VIII administered is expressed in International Units (IU), which are related to the current WHO standard for factor VIII products. The activity of factor VIII in plasma is expressed either as percentage (relative to normal level human plasma) or in International Units (relative to an International Standard for factor VIII in plasma).

One International Unit (IU) of factor VIII activity is equivalent to that quantity of factor VIII in one ml normal human plasma.

On-demand treatment

The calculation of the required dose of factor VIII is based on the empirical finding that 1 International Unit (IU) factor VIII per kg body weight raises the plasma factor VIII activity by 2 IU/dl. The required dose is determined using the following formula:

Required units = body weight (kg) x desired factor VIII rise (%) (IU/dl) x 0.5 (IU/kg per IU/dl).

The amount to be administered and the frequency of administration should always be oriented to the clinical effectiveness in the individual case.

In the case of the following haemorrhagic events, the factor VIII activity should not fall below the given plasma activity level (in % of normal or IU/dl) in the corresponding period. The following table can be used to guide dosing in bleeding episodes and surgery:

Table 1 Guide for dosing in bleeding episodes and surgery

Degree of haemorrhage/Type of surgical procedure	FVIII level required (%) (IU/dl)	Frequency of doses (hours)/Duration of therapy (days)
<u>Haemorrhage</u>		
Early haemarthrosis, muscle bleeding or oral bleeding	20–40	Repeat every 12 to 24 hours, at least 1 day, until the bleeding episode as indicated by pain is resolved or healing achieved
More extensive haemarthrosis, muscle bleeding or haematoma	30–60	Repeat infusion every 12–24 hours for 3–4 days or more until pain and acute disability are resolved

Life threatening haemorrhages	60–100	Repeat infusion every 8 to 24 hours until threat is resolved
<hr/>		
<u>Surgery</u>		
<i>Minor surgery including tooth extraction</i>	30–60	Every 24 hours, at least 1 day, until healing is achieved
<i>Major surgery</i>	80–100 (pre- and postoperative)	Repeat infusion every 8–24 hours until adequate wound healing, then therapy for at least another 7 days to maintain a factor VIII activity of 30% to 60% (IU/dl)
<hr/>		

Prophylaxis

For long term prophylaxis against bleeding in patients with severe haemophilia A. The usual recommended doses are 20–40 IU of factor VIII per kg body weight every second day or 20–50 IU of factor VIII per kg body weight 3 times weekly. In adults and adolescents (>12 years) a less frequent regimen (40–60 IU/kg every third day or twice weekly) may be applicable. In some cases, especially in younger patients, shorter dosage intervals or higher doses may be necessary.

Treatment monitoring

During the course of treatment, appropriate determination of factor VIII levels is advised to guide the dose to be administered and the frequency of repeated injections. In the case of major surgical interventions in particular, precise monitoring of the substitution therapy by means of coagulation analysis (plasma factor VIII activity) is indispensable. Individual patients may vary in their response to factor VIII, demonstrating different half-lives and recoveries. Dose based on bodyweight may require adjustment in underweight or overweight patients. When using an *in vitro* thromboplastin time (aPTT) based one stage clotting assay for determining factor VIII activity in patients' blood samples, plasma factor VIII activity results can be significantly affected by both the type of aPTT reagent and the reference standard used in the assay. Also there can be significant discrepancies between assay results obtained by aPTT-based one stage clotting assay and the chromogenic assay according to Ph. Eur. This is of importance particularly when changing the laboratory and/or reagents used in the assay.

Surgery

There is limited experience of surgery in paediatric patients.

Elderly

There is no experience in patients > 65 years.

Paediatric population

For long term prophylaxis against bleeding in patients below the age of 12, doses of 25–50 IU of factor VIII per kg body weight every second day or 25–60 IU of factor VIII per kg body weight 3 times weekly are recommended. For paediatric patients above the age of 12 the dose recommendations are the same as for adults.

Method of administration

Intravenous use.

The recommended infusion rate for NovoEight is 1–2 ml/min. The rate should be determined by the patient's comfort level.

For instructions on reconstitution of the medicinal product before administration, see section 6.6.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Known allergic reaction to hamster proteins.

4.4 Special warnings and precautions for use

Hypersensitivity

Allergic type hypersensitivity reactions are possible with NovoEight. The product contains traces of hamster proteins, which in some patients may cause allergic reactions. If symptoms of hypersensitivity occur, patients should be advised to discontinue use of the medicinal product immediately and contact their physician. Patients should be informed of the early signs of hypersensitivity reactions including hives, generalised urticaria, tightness of the chest, wheezing, hypotension and anaphylaxis.

In case of shock, standard medical treatment for shock should be implemented.

Inhibitors

The formation of neutralising antibodies (inhibitors) to factor VIII is a known complication in the management of individuals with haemophilia A. These inhibitors are usually IgG immunoglobulins directed against the factor VIII procoagulant activity, which are quantified in Bethesda Units (BU) per ml of plasma using the modified assay. The risk of developing inhibitors is correlated to the severity of the disease as well as the exposure to factor VIII, this risk being highest within the first 20 exposure days. Rarely, inhibitors may develop after the first 100 exposure days.

Cases of recurrent inhibitor (low titre) have been observed after switching from one factor VIII product to another in previously treated patients with more than 100 exposure days who have a previous history of inhibitor development. Therefore, it is recommended to monitor all patients carefully for inhibitor occurrence following any product switch.

The clinical relevance of inhibitor development will depend on the titre of the inhibitor, with low titre inhibitors which are transiently present or remain consistently low titre posing less of a risk of insufficient clinical response than high titre inhibitors.

In general, all patients treated with coagulation factor VIII products should be carefully monitored for the development of inhibitors by appropriate clinical observation and laboratory test. If the expected factor VIII activity plasma levels are not attained, or if bleeding is not controlled with an appropriate dose, testing for factor VIII inhibitor presence should be performed. In patients with high levels of inhibitor, factor VIII therapy may not be effective and other therapeutic options should be considered. Management of such patients should be directed by physicians with experience in the care of haemophilia and factor VIII inhibitors.

It is strongly recommended that every time that NovoEight is administered to a patient, the name and batch number of the product are recorded in order to maintain a link between the patient and the batch of the medicinal product.

Excipient related considerations

After reconstitution this medicinal product contains 0.31 mmol sodium (7 mg) per ml of reconstituted solution. To be taken into consideration by patients on a controlled sodium diet.

Cardiovascular event

In patients with existing cardiovascular risk factors, substitution therapy with FVIII may increase the cardiovascular risk.

Catheter-related complications

If a central venous access device (CVAD) is required, risk of CVAD-related complications including local infections, bacteraemia and catheter site thrombosis should be considered.

Paediatric population

The listed warnings and precautions apply both to adults and children.

4.5 Interaction with other medicinal products and other forms of interaction

No interaction studies have been performed with NovoEight.

4.6 Fertility, pregnancy and lactation

Animal reproduction studies have not been conducted with NovoEight. Based on the rare occurrence of haemophilia A in women, experience regarding the use of factor VIII during pregnancy and breastfeeding is not available. Therefore, factor VIII should be used during pregnancy and breastfeeding only if clearly indicated.

4.7 Effects on ability to drive and use machines

NovoEight has no influence on the ability to drive and use machines.

4.8 Undesirable effects

Summary of the safety profile

Hypersensitivity or allergic reactions (which may include angioedema, burning and stinging at the infusion site, chills, flushing, generalised urticaria, headache, hives, hypotension, lethargy, nausea, restlessness, tachycardia, tightness of the chest, tingling, vomiting, wheezing) have been observed rarely and may in some cases progress to severe anaphylaxis (including shock).

Very rarely development of antibodies to hamster protein with related hypersensitivity reactions has been observed.

Development of neutralising antibodies (inhibitors) may occur in patients with haemophilia A treated with factor VIII, including with NovoEight, see section 5.1. If such inhibitors occur, the condition will manifest itself as an insufficient clinical response. In such cases, it is recommended that a specialised haemophilia centre is contacted.

Tabulated list of adverse reactions

The table presented below is according to the MedDRA system organ classification (SOC and Preferred Term Level).

Frequencies have been evaluated according to the following convention: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$), and not known (cannot be estimated from the available data).

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Table 2 Frequency of adverse drug reactions in clinical trials

System Organ Class	Frequency* in PTPs	Frequency* in PUPs	Adverse reaction
Blood and lymphatic system disorders	Uncommon**	Very common**	FVIII inhibition
Psychiatric disorders	Uncommon		Insomnia
Nervous system disorders	Uncommon		Headache, dizziness, burning sensation
Cardiac disorders	Uncommon		Sinus tachycardia, acute myocardial infarction
Vascular disorders	Uncommon		Hypertension, lymphoedema, hyperaemia
		Common	Flushing, Thrombophlebitis superficial
Skin and subcutaneous tissue disorders		Common	Rash, rash erythematous
	Uncommon		Rash, lichenoid keratosis, skin burning sensation
Musculoskeletal and connective tissue disorders	Uncommon		Musculoskeletal stiffness, arthropathy, pain in extremity, musculoskeletal pain
		Common	Haemarthrosis, Muscle haemorrhage
Respiratory, thoracic and mediastinal disorders		Common	Cough
General disorders and administration site conditions	Common		Injection site reactions***
		Common	Pyrexia, catheter site erythema
	Uncommon		Fatigue, feeling hot, oedema peripheral, pyrexia
Investigations	Common		Hepatic enzymes increased****
		Common	Anti factor VIII antibody positive
	Uncommon		Heart rate increased
Gastrointestinal disorders		Common	Vomiting
Injury, poisoning and procedural complications	Common		Incorrect dose administered
		Common	Infusion related reaction
	Uncommon		Contusion
Product issues		Common	Thrombosis in device

* Calculated based on total number of unique patients in all clinical trials (301), of which 242 were previously treated patients (PTPs) and 59 were previously untreated patients (PUPs).

** Frequency is based on studies with all FVIII products which included patients with severe haemophilia A.

*** Injection site reactions include injection site erythema, injection site extravasation and injection site pruritus.

**** *Hepatic enzymes increased include alanine aminotransferase, aspartate aminotransferase, gamma-glutamyltransferase and bilirubin.*

Description of selected adverse reactions

During all clinical studies with NovoEight in previously treated patients, a total of 35 adverse reactions were reported in 23 of 242 patients exposed to NovoEight. The most frequently reported adverse reactions were injection site reactions, incorrect dose administered and hepatic enzymes increased. Of the 35 adverse reactions, 2 were reported in 1 out of 31 patients below 6 years of age, none in patients from 6 to \leq 12 years of age, 1 event in 1 out of 24 patients (12 to < 18 years of age) and 32 were reported in 21 out of 155 adults (\geq 18 years).

Paediatric population

In clinical trials involving 63 previously treated paediatric patients between 0 and 12 years of age and 24 adolescents between 12 and 18 years of age with severe haemophilia A no difference in the safety profile of NovoEight was observed between paediatric patients and adults.

In the trial with previously untreated patients, between 0 and 6 years of age, a total of 36 adverse reactions were reported in 32 of 59 patients exposed to NovoEight. The most frequently reported adverse reaction was Factor VIII inhibition, see section 4.4. High risk genetic mutations were identified in 91.7% of the overall and 93.3% of the high titre inhibitors. No other factors were significantly associated with inhibitor development.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

4.9 Overdose

No symptoms of overdose with recombinant coagulation factor VIII have been reported.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antihemorrhagics, blood coagulation factor VIII, ATC code: B02BD02.

Mechanism of action

NovoEight contains turoctocog alfa, a human coagulation factor VIII (rDNA), with a truncated B-domain. This glycoprotein has the same structure as human factor VIII when activated, and post-translational modifications that are similar to those of the plasma-derived molecule. The tyrosine sulphation site present at Tyr1680 (native full length), which is important for the binding to von Willebrand factor, has been found to be fully sulphated in the turoctocog alfa molecule. When infused into a haemophilia patient, factor VIII binds to endogenous von Willebrand Factor in the patient's circulation. The factor VIII/von Willebrand factor complex consists of two molecules (factor VIII and von Willebrand factor) with different physiological functions. Activated factor VIII acts as a co-factor for activated factor IX, accelerating the conversion of factor X to activated factor X. Activated factor X converts prothrombin into thrombin. Thrombin then converts fibrinogen into fibrin and a clot can be formed. Haemophilia A is a sex-linked hereditary disorder of blood coagulation due to decreased levels of factor VIII:C and results in profuse bleeding into joints, muscles or internal organs, either spontaneously or as a result of accidental or surgical trauma. By replacement therapy the plasma levels of factor VIII are increased, thereby enabling a temporary correction of the factor deficiency and correction of bleeding tendencies.

Clinical efficacy

Four multi-centre, open-labelled, non-controlled trials have been conducted to evaluate the safety and efficacy of NovoEight in the prevention and treatment of bleeds and during surgery in patients with severe haemophilia A (FVIII activity $\leq 1\%$). Three of these trials were performed in previously treated patients and the fourth in previously untreated patients. The trials included 297 exposed patients; 175 adolescents or adult patients without inhibitors from the age of 12 years (≥ 150 exposure days), 63 previously treated paediatric patients without inhibitors below 12 years of age (≥ 50 exposure days) and 59 previously untreated patients below 6 years of age.

188 out of 238 previously treated patients continued into the safety extension trial. Treatment with NovoEight was shown to be safe and had the intended haemostatic and preventive effect.

Of the 3,153 reported bleeds observed in 297 of the patients, 2,794 (88.6%) of the bleeds were resolved with 1-2 infusions of NovoEight.

Table 3 Consumption of turoctocog alfa and overall success rates in previously treated patients

	Younger children (0 – < 6 years)	Older children (6 – < 12 years)	Adolescents (12 – < 18 years)	Adults (≥ 18 years)	Total
Number of patients	31	32	24	151	238
Dose used for prevention per patient (IU/kg BW)					
Mean (SD)	41.5 (8.1)	38.4 (9.4)	28.5 (9.3)	28.5 (8.3)	31.9 (10.1)
Min ; Max	3.4 ; 196.3	3.2 ; 62.5	17.4 ; 73.9	12.0 ; 97.4	3.2 ; 196.3
Dose used for treatment of bleed (IU/kg BW)					
Mean (SD)	44.0 (12.6)	40.4 (10.5)	29.3 (10.3)	35.0 (12.3)	36.0 (12.5)
Min ; Max	21.4 ; 193.8	24.0 ; 71.4	12.4 ; 76.8	6.4 ; 104.0	6.4 ; 193.8
Success rate* %	92.2%	88.4%	85.1%	89.6%	89.2%

BW: Body weight, SD: Standard deviation

*Success is defined as either 'Excellent' or 'Good'.

A total of 30 surgeries were performed in 25 patients of which 26 were major surgeries and 4 were minor. Haemostasis was successful in all surgeries and no treatment failures were reported.

5.2 Pharmacokinetic properties

All pharmacokinetic studies with turoctocog alfa were conducted in previously treated patients with severe haemophilia A (FVIII $\leq 1\%$). The analysis of plasma samples was conducted using both the one-stage clotting assay and the chromogenic assay.

In an international study involving 36 laboratories, the assay performance of NovoEight in FVIII:C assays was evaluated and compared to a marketed full length recombinant FVIII product. The study showed that comparable and consistent results were obtained for both products and that NovoEight can be reliably measured in plasma without the need of a separate NovoEight standard.

The single dose pharmacokinetic parameters of NovoEight are listed in Table 4 for the clotting assay and in Table 5 for the chromogenic assay.

Table 4 Single-dose pharmacokinetics of turoctocog alfa in patients with severe haemophilia A (FVIII \leq 1%), clotting assay

Parameter	0 – < 6 years	6 – < 12 years	\geq 12 years
	n=14	n=14	n=33
	Mean (SD)	Mean (SD)	Mean (SD)
Incremental recovery (IU/ml)/(IU/kg)	0.018 (0.007)	0.020 (0.004)	0.022 (0.004)
AUC ((IU*h)/ml)	9.92 (4.11)	11.09 (3.74)	15.26 (5.77)
CL (ml/h/kg)	6.21 (3.66)	5.02 (1.68)	3.63 (1.09)
$t_{1/2}$ (h)	7.65 (1.84)	8.02 (1.89)	11.00 (4.65)
V_{ss} (ml/kg)	56.68 (26.43)	46.82 (10.63)	47.40 (9.21)
C_{max} (IU/ml)	1.00 (0.58)	1.07 (0.35)	1.226 (0.41)
Mean residence time (h)	9.63 (2.50)	9.91 (2.57)	14.19 (5.08)

Table 5 Single-dose pharmacokinetics of turoctocog alfa in patients with severe haemophilia A (FVIII \leq 1%), chromogenic assay

Parameter	0 – < 6 years	6 – < 12 years	\geq 12 years
	n=14	n=14	n=33
	Mean (SD)	Mean (SD)	Mean (SD)
Incremental recovery (IU/ml)/(IU/kg)	0.022 (0.006)	0.025 (0.006)	0.029 (0.006)
AUC ((IU*h)/ml)	12.23 (4.36)	14.37 (3.48)	19.63 (7.73)
CL (ml/h/kg)	4.59 (1.73)	3.70 (1.00)	2.86 (0.94)
$t_{1/2}$ (h)	9.99 (1.71)	9.42 (1.52)	11.22 (6.86)
V_{ss} (ml/kg)	55.46 (23.53)	41.23 (6.00)	38.18 (10.24)
C_{max} (IU/ml)	1.12 (0.31)	1.25 (0.27)	1.63 (0.50)
Mean residence time (h)	12.06 (1.90)	11.61 (2.32)	14.54 (5.77)

The pharmacokinetic parameters were comparable between paediatric patients below 6 years of age and the paediatric patients from 6 to below 12 years of age. Some variation was observed in the pharmacokinetic parameters of NovoEight between paediatric and adult patients. The higher CL and the shorter $t_{1/2}$ seen in paediatric patients compared to adult patients with haemophilia A may be due in part to the known higher plasma volume per kilogram body weight in younger patients.

5.3 Preclinical safety data

Non-clinical data reveal no special concern for humans based on conventional studies of safety pharmacology and repeated dose toxicity.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Powder:

Sodium chloride
L-histidine
Sucrose
Polysorbate 80
L-methionine
Calcium chloride dihydrate
Sodium hydroxide
Hydrochloric acid

Solvent:

Sodium chloride
Water for injections

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

Unopened vial

30 months when stored in a refrigerator (2°C – 8°C).

During the shelf life, the product may be kept at:

- room temperature ($\leq 30^{\circ}\text{C}$) for a single period no longer than 9 months
- **or**
- above room temperature (30°C up to 40°C) for a single period no longer than 3 months.

Once the product has been taken out of the refrigerator, the product must not be returned to the refrigerator.

Please record the beginning of storage and the storage temperature on the product carton.

After reconstitution:

Chemical and physical in-use stability have been demonstrated for:

- 24 hours stored at $2^{\circ}\text{C} - 8^{\circ}\text{C}$
- 4 hours stored at 30°C , for product which has been kept for a single period no longer than 9 months at room temperature ($\leq 30^{\circ}\text{C}$)
- 4 hours stored up to 40°C , for product which has been kept for a single period no longer than 3 months at above room temperature (30°C up to 40°C)

From a microbiological point of view, the medicinal product should be used immediately after reconstitution. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than as stated above, unless reconstitution has taken place in controlled and validated aseptic conditions.

Any unused reconstituted product stored at room temperature ($\leq 30^{\circ}\text{C}$) or up to 40°C for more than 4 hours should be discarded.

6.4 Special precautions for storage

Store in refrigerator ($2^{\circ}\text{C} - 8^{\circ}\text{C}$).

Do not freeze.

Keep the vial in the outer carton in order to protect from light.

For storage at room temperature ($\leq 30^{\circ}\text{C}$) or up to 40°C and storage conditions after reconstitution of the medicinal product, see section 6.3.

6.5 Nature and contents of container

Each pack of NovoEight 250 IU, 500 IU powder and solvent for solution for injection contains:

- 1 glass vial (type I) with powder and chlorobutyl rubber stopper
- 1 sterile vial adapter for reconstitution
- 1 pre-filled syringe of 4 ml solvent with backstop (polypropylene), a rubber plunger (bromobutyl) and a syringe cap with a stopper (bromobutyl)
- 1 plunger rod (polypropylene).

6.6 Special precautions for disposal and other handling

NovoEight is to be administered intravenously after reconstitution of the powder with the solvent supplied in the syringe. After reconstitution the solution appears as a clear or slightly opalescent solution. Do not use solutions that are cloudy or have deposits.

You will also need an infusion set (tubing and butterfly needle), sterile alcohol swabs, gauze pads and plasters. These devices are not included in the NovoEight package.

Always use an aseptic technique.

Reconstitution

A)

Take the vial, the vial adapter and the pre-filled syringe out of the carton. Leave the plunger rod untouched in the carton. Bring the vial and the pre-filled syringe to room temperature. You can do this by holding them in your hands until they feel as warm as your hands. Do not use any other way to heat the vial and pre-filled syringe.



B)

Remove the plastic cap from the vial. If the plastic cap is loose or missing, do not use the vial. Wipe the rubber stopper on the vial with a sterile alcohol swab and allow it to air dry for a few seconds before use.



C)

Remove the protective paper from the vial adapter. If the protective paper is not fully sealed or if it is broken, do not use the vial adapter. Do not take the vial adapter out of the protective cap with your fingers.




D)

Turn over the protective cap and snap the vial adapter onto the vial. Once attached do not remove the vial adapter from the vial.




E)

Lightly squeeze the protective cap with your thumb and index finger as shown. Remove the protective cap from the vial adapter.



F)

Grasp the plunger rod by the wide top and immediately connect the plunger rod to the syringe by turning it clockwise into the plunger inside the pre-filled syringe until resistance is felt.




G)

Remove the syringe cap from the pre-filled syringe by bending it down until the perforation breaks. Do not touch the syringe tip under the syringe cap.



H)

Screw the pre-filled syringe securely onto the vial adapter until resistance is felt.



I)

Hold the pre-filled syringe slightly tilted with the vial pointing downwards. Push the plunger rod to inject all the solvent into the vial.



J)

Keep the plunger rod pressed down and swirl the vial gently until all the powder is dissolved. Do not shake the vial as this will cause foaming.



It is recommended to use NovoEight immediately after reconstitution. For storage conditions of the reconstituted medicinal product see section 6.3.

If a larger dose is needed, repeat steps A to J with additional vials, vial adapters and pre-filled syringes.

Administration of the reconstituted solution

K)

Keep the plunger rod pushed completely in.
Turn the syringe with the vial upside down. Stop pushing the plunger rod and let it move back on its own while the reconstituted solution fills the syringe. Pull the plunger rod slightly downwards to draw the reconstituted solution into the syringe.


In case you only need part of the entire vial, use the scale on the syringe to see how much reconstituted solution you withdraw, as instructed by your doctor or nurse.

While holding the vial upside down, tap the syringe gently to let any air bubbles rise to the top. Push the plunger rod slowly until all air bubbles are gone.



L)

Unscrew the vial adapter with the vial.



NovoEight is now ready for injection. Locate a suitable site and slowly inject NovoEight into the vein over a period of 2-5 minutes.

Disposal

After injection, safely dispose of all unused NovoEight solution, the syringe with the infusion set, the vial with the vial adapter and other waste materials as instructed by your pharmacist.

Do not throw it out with the ordinary household waste.

7. MARKETING AUTHORISATION HOLDER

Novo Nordisk Pharma (Thailand) Ltd., Bangkok, Thailand

8. MARKETING AUTHORISATION NUMBERS

NovoEight 250 IU
1C 43/60 (NBC)

NovoEight 500 IU
1C 42/60 (NBC)

9. DATE OF FIRST AUTHORISATION

07 December 2017

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the Thai FDA.