

เอกสารกำกับยาภาษาอังกฤษ

Leaflet

MAGTEN

Each 10 mL contains:-

Magnesium Sulfate 10% ๑๐/๑๐ ๑๕/๑๒/๑๕

Indication : Treatment and prevention of Hypomagnesemia and in seizure, prevention in severe preclampsia or eclampsia, central depressant in convulsions.

Dosage :

Hypomagnesemia

- Neonates :- 25-50 mg/kg/dose IV every 8-12 hours for 2-3 doses.
- Children :- 25-50 mg/kg/dose IM or IV every 4-6 hours for 3-4 doses.
Maximum single dose 2,000 mg. may repeat if Hypomagnesemia persists, maintenance 30-60 mg/kg/day
- Adult :- 1 g. every 6 hours IM or IV for 4 doses, for severe Hypomagnesemia 8-12 g. MgSO₄/day in divided has been used.

Management of Seizures and Hypertension

- Children :- 20-100 mg/kg/dose IM or IV every 4-6 hours as needed. in severe cases doses as high as 200 mg/kg/dose have been used.

Eclampsia, Pre-eclampsia

- Adult :- I.M. 1-4 g. every 4 hours
I.V. Initial 4 g., then 1-4 g/hour by continuous infusion maximum dose should not exceed 30-40 g/day, maximum rate of infusion : 1-2 g/hour.

Warnings/Precaution : Use with caution in patients with impaired renal function, Use with caution in digitalized patients. The following may be especially important in patient monitering : blood pressure, ECG, serum magnesium concentration, respiratory rate, deep tendon reflex, renal function when MgSO₄ is administered parenterally.

Adverse Reaction : Hypermagnesemia.

Drug Interaction : Enhances neuromuscular blockers, digitalis glycosides.

Manufactured by : UMEDA CO., LTD.
26/18-21 RAM-INDRA ROAD,
BANGKHEN, BANGKOK, THAILAND.
Tel. (662) 943-5588 Fax. (662)943-5038