

## Summary of Product Characteristics

### 1. Name of the Medicinal Product

Kolimycin, 150 mg, Powder for solution for injection, infusion or inhalation.

### 2. Quality and Quantitative Composition

Each vial contains Colistimethate Sodium eq. to Colistin 150 mg.

### 3. Pharmaceutical Form

Powder for solution for injection, infusion or inhalation.

White to yellowish white sterile powder.

### 4. Clinical Particulars

#### 4.1 Therapeutic indication:

**Kolimycin by intravenous administration** is indicated in adults and children including neonates for the treatment of serious infections due to selected aerobic Gram-negative pathogens in patients with limited treatment options (see sections 4.2, 4.4, 4.8 and 5.1).

**Kolimycin by inhalation** is also indicated for the management of adult and paediatric chronic pulmonary infections due to *Pseudomonas aeruginosa* in patients with cystic fibrosis (see section 5.1).

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

#### 4.2 Posology and method of administration:

##### SYSTEMIC TREATMENT

The dose to be administered and the treatment duration should take into account the severity of the infection as well as the clinical response. Therapeutic guidelines should be adhered to.

The dose is expressed in IU of colistimethate sodium (CMS). A conversion table from CMS in IU to mg of CMS as well as to mg of colistin base activity (CBA) is included at the end of this section.

##### Posology

The following dose recommendations are made based on limited population-pharmacokinetic data in critically ill patients (see section 4.4):

##### *Adults and adolescents*

- Maintenance dose 9 MIU (300 mg of CBA) /day in 2-3 divided doses.
- In patients who are critically ill, a loading dose of 9 MIU (300 mg of CBA) should be administered.
- The most appropriate time interval to the first maintenance dose has not been established.
- Modelling suggests that loading and maintenance doses of up to 12 MIU (400 mg of CBA) may be required in patients with good renal function in some cases. Clinical experience with such doses is however extremely limited, and safety has not been established.

- The loading dose applies to patients with normal and impaired renal functions including those on renal replacement therapy.

*Renal impairment*

- Dose adjustments in renal impairment are necessary, but pharmacokinetic data available for patients with impaired renal function is very limited.
- The following dose adjustments are suggested as guidance.
- Dose reductions are recommended for patients with creatinine clearance < 50 ml/min:
- Twice daily dosing is recommended.

| Creatinine clearance (ml/min) | Daily dose (MIU) | Daily dose (mg of CBA) |
|-------------------------------|------------------|------------------------|
| < 50-30                       | 5.5-7.5 MIU      | 183-250 mg.            |
| < 30-10                       | 4.5-5.5 MIU      | 150-183 mg.            |
| < 10                          | 3.5 MIU          | 116 mg.                |

MIU = million IU, 1 mg = 30,000 IU, CBA = Colistin base activity

- Haemodialysis and continuous haemo(dia)filtration

Colistin appears to be dialyzable through conventional haemodialysis and continuous venovenous haemo(dia)filtration (CVVHF, CVVHDF). There are extremely limited data from population PK studies from very small numbers of patients on renal replacement therapy. Firm dose recommendations cannot be made. The following regimes could be considered.

- Haemodialysis

No-HD days: 2.25 MIU (75 mg of CBA)/day (2.2-2.3 MIU (73-77 mg of CBA) /day)

HD days: 3 MIU (100 mg of CBA) /day on haemodialysis day, to be given after the HD session.

Twice daily dosing is recommended.

- CVVHF / CVVHDF

As in patients with normal renal function. Three times daily dosing is recommended.

*Hepatic impairment*

There are no data in patients with hepatic impairment. Caution is advised when administering colistimethate sodium in these patients.

*Elderly*

No dose adjustments in older patients with normal renal function are considered necessary.

*Paediatric population*

The data supporting the dose regimen in paediatric are very limited. Renal maturity should be taken into consideration when selecting the dose. The dose should be based on lean body weight.

- Children < 40kg
  - 75,000-150,000 IU (2.5-5 mg of CBA) /kg/day divided into 3 doses.
  - For children with a body weight above 40 kg, use of the dosing recommendation for adults should be considered.
  - The use of doses > 150,000 IU (5 mg of CBA) /kg/day has been reported in children with cystic fibrosis.
  - There are no data regarding the use or magnitude of a loading dose in critically ill children.
  - No dose recommendations have been established in children with impaired renal function.

*Intrathecal and intraventricular administration*

- Based on limited data, the following dose is recommended in adults: Intraventricular route: 125,000 (0.4 mg of CBA) /day
- Intrathecally administered doses should not exceed those recommended for intraventricular use.
- No specific dosing recommendation can be made in children for intrathecal and intraventricular routes of administration.

Method of administration

Kolimycin is administered intravenously as a slow infusion over 30-60 minutes.

Patients with a totally implantable venous access device (TIVAD) in place may tolerate a bolus injection of up to 2 MIU (68 mg of CBA) in 10ml given over a minimum of 5 minutes (see section 6.6).

Colistimethate sodium undergoes hydrolysis to the active substance colistin in aqueous solution. For dose preparation, particularly where combination of multiple vials is needed, reconstitution of the required dose must be performed using strict aseptic technique (see section 6.6).

**Dose conversion table:**

In the EU, the dose of colistimethate sodium (CMS) must be prescribed and administered only as IU. However, the product label stated the number of mg per vial.

Confusion and medication errors have occurred because of the different expressions of dose in terms of potency. The dose is expressed in the US, and other parts of the world, as milligrams of colistin base activity (mg CBA).

The following conversion table is prepared for information and the values must be considered nominal and approximate only.

## CMS conversion table

| Potency   |                             | $\approx$ mg CMS |
|-----------|-----------------------------|------------------|
| IU        | $\approx$ mass of CBA (mg)* |                  |
| 12 500    | 0.4                         | 1                |
| 150 000   | 5                           | 12               |
| 1 000 000 | 34                          | 80               |
| 4 500 000 | 150                         | 360              |
| 9 000 000 | 300                         | 720              |

\* Nominal potency of the drug substance = 12,500 IU/mg

## AEROSOL INHALATION

It is recommended that colistimethate sodium (CMS) should be administered under the supervision of physicians with appropriate experience in its use.

### Posology

The dosage can be adjusted depending on the severity of the condition and clinical response.

Recommended dose range:

### **Administration via inhalation**

*Adults, adolescents and children  $\geq$  2 years*

1-2 MIU (34-68 mg of CBA) two to three times per day (max 6 MIU (200 mg of CBA) / day)

*Children < 2 years*

0.5-1 MIU (17-34 mg of CBA) twice daily (max 2 MIU (68 mg of CBA) / day)

Relevant clinical guidance on treatment regimens, including duration of treatment. Periodicity and co-administration of other antibacterial agents should be adhered to.

*Elderly*

Dose adjustment is not considered necessary.

*Renal impairment*

Dose adjustment is not considered necessary; however, caution is advised in patients with renal impairment (see sections 4.4 and 5.2).

*Hepatic impairment*

Dose adjustment is not considered necessary.

## Method of administration

### **For inhalation use.**

Colistimethate sodium undergoes hydrolysis to the active substance colistin in aqueous solution. For special precautions for disposal and handling of reconstituted solutions, see section 6.6.

If other treatments are being taken, they should be taken in the order recommended by the physician.

See above for Dose conversion table.

### **4.3 Contraindications**

Hypersensitivity to colistimethate sodium (colistin) or to polymyxin B.

### **4.4 Special warnings and precautions for use**

- Consideration should be given to co-administering intravenous colistimethate sodium with another antibacterial agent whenever this is possible, taking into account the remaining susceptibilities of the pathogen(s) under treatment. As the development of resistance to intravenous colistin has been reported in particular when it is used as a monotherapy, co-administration with other antibacterial should also be considered in order to prevent the emergence of resistance.
- There are limited clinical data on the efficacy and safety of intravenous colistimethate sodium. The recommended doses in all subpopulations are equally based on limited data (clinical and pharmacokinetic/ pharmacodynamics data). In particular there are limited safety data for the use of high doses ( $> 6$  MIU (200 mg of CBA) / day) and the use of a loading dose, and for special populations (patients with renal impairment and the paediatric population). Colistimethate sodium should only be used when other, more commonly prescribed antibiotics are not effective or not appropriate.
- Renal function monitoring should be performed at the start of treatment and regularly during treatment in all patients. The dose of colistimethate sodium should be adjusted according to creatinine clearance (see section 4.2). Patients who are hypovolaemic or those potentially nephrotoxic drugs are at increased risk of nephrotoxicity from colistin (see section 4.5 and 4.8). Nephrotoxicity has been reported to be associated with cumulative dose and treatment duration in some studies. The benefit of prolonged treatment duration should be balanced against the potentially increase risk of renal toxicity.
- Caution is advised when administering colistimethate sodium to infants  $< 1$  year of age as renal function is not fully mature in this age group. Further, the effect of immature renal and metabolic function on the conversion of colistimethate sodium to colistin is not known.
- In case of an allergic reaction, treatment with colistimethate sodium must be discontinued and appropriate measures implemented.
- High serum concentrations of colistimethate sodium, which may be associated with overdosage or failure to reduce the dosage in patients with renal impairment, have been reported to lead to neurotoxic effects such as facial paraesthesia, muscle weakness, vertigo, slurred speech, vasomotor instability, visual disturbances, confusion, psychosis and apnoea.

- Monitoring should be performed for perioral paraesthesia and paraesthesia in the extremities, which are signs of overdose (see section 4.9).
- Colistimethate sodium is known to reduce the presynaptic release of acetyl-choline at the neuro-muscular junction and should be used in patients with myasthenia gravis with the greatest caution and only if clearly needed.
- Respiratory arrest has been reported following intramuscular administration of colistimethate sodium. Impaired renal function increases the possibility of apnoea and neuromuscular blockade following administration of colistimethate sodium.
- Colistimethate sodium should be used with extreme caution in patients with porphyria.
- Antibiotic-associated colitis and pseudomembranous colitis have been reported with nearly all anti-bacterial agents and may occur with colistimethate sodium. They may range from mild to life-threatening in severity. It is important to consider this diagnosis in patients who develop diarrhoea during or after the use of colistimethate sodium (see section 4.8).
- Discontinuation of therapy and the administration of specific treatment for *Clostridium difficile* should be considered.
- Medicinal products that inhibit peristalsis should not be given.
- Intravenous colistimethate sodium does not cross the blood brain barrier to a clinically relevant extent. The use of intrathecal or intraventricular administration of colistimethate sodium in the treatment of meningitis was not systematically investigated in clinical trials and is supported by case reports only. Data supported the posology are very limited. The most commonly observed adverse effect of CMS administration was aseptic meningitis (see section 4.8).
- Bronchospasm may occur on inhalation of antibiotics. This may be prevented or treated with appropriate use of beta2-agonists. If troublesome, treatment should be withdrawn.

#### 4.5 Interaction with other medicinal products and other forms of interaction

- Concomitant use of intravenous colistimethate sodium with other medications that are potentially nephrotoxic or neurotoxic should be undertaken with great caution.
- Caution should be taken with concomitant use with other formulations of colistimethate sodium as there is little experience and there is a possibility of summative toxicity.
- No. in vivo interaction studies have been performed. The mechanism of conversion of colistimethate sodium to the active substance, colistin, is not characterized. The mechanism of colistin clearance, including renal handling, is equally unknown. Colistimethate sodium or colistin did not induce the activity of any P 450 (CYP) enzyme tested (CYP1A2, 2B6, 2C8, 2C9, 2C19 and 3A4/5) in in vitro studies in human hepatocytes.
- The potential for drug-drug interactions should be borne in mind when Kolimycin is co-administered with drugs known to inhibit or induce drug metabolizing enzymes or drugs known to be substrates for renal carrier mechanisms.
- Due to the effects of colistin on the release of acetylcholine, non-depolarising muscle relaxants should be used with caution in patients receiving colistimethate sodium as their effects could be prolonged (see section 4.4).

- Co-treatment with colistimethate sodium and macrolides such as azithromycin and clarithromycin or fluoroquinolones such as norfloxacin and ciprofloxacin should be undertaken with caution in patients with myasthenia gravis (see section 4.4).
- Concomitant use of colistimethate sodium with other medicinal products of neurotoxic and/or nephrotoxic potential should be avoided. These include the aminoglycoside antibiotics such as gentamicin, amikacin, netilmicin and tobramycin. There may be an increased risk of nephrotoxicity if given concomitantly with cephalosporin antibiotics.

#### 4.6 Pregnancy and lactation

There are no adequate data from the use of colistimethate sodium in pregnant women. Single dose studies in human pregnancy show that colistimethate sodium crosses the placental barrier and there may be a risk of foetal toxicity if repeated doses are given to pregnant patients. Animal studies are insufficient with respect to the effect of colistimethate sodium on reproduction and development (see Section 5.3-Preclinical safety data). Colistimethate sodium should be used in pregnancy only if the benefit to the mother outweighs the potential risk to the fetus.

Colistimethate sodium is secreted in breast milk. Colistimethate sodium should be administered to breastfeeding women only when clearly needed.

#### 4.7 Effects on ability to drive and use machines

During parenteral treatment with colistimethate sodium neurotoxicity may occur with the possibility of dizziness, confusing or visual disturbance. Patients should be warned not to drive or operate machinery if these effects occur.

#### 4.8 Undesirable effects

##### Systemic treatment

- The likelihood of adverse events may be related to the age, renal function and condition of the patient.
- In cystic fibrosis patients neurological events have been reported in up to 27% of patients. These are generally mild and resolve during or shortly after treatment.
- Neurotoxicity may be associated with overdose, failure to reduce the dose in patients with renal insufficiency and concomitant use of either neuromuscular blocking drugs or other drugs with similar neurological effects. Reducing the dose may alleviate symptoms. Effects may include apnoea, transient sensory disturbances (such as facial paraesthesia and vertigo) and, rarely, vasomotor instability, slurred speech, visual disturbances, confusion or psychosis.
- Adverse effects on renal function have been reported, usually following use of higher than recommended doses in patients with normal renal function, or failure to reduce the dosage in patients with renal impairment or during concomitant use of other nephrotoxic drugs. The effects are usually reversible on discontinuation of therapy.
- In cystic fibrosis patients treated within the recommended dosage limits, nephrotoxicity appears to be rare (less than 1%). In seriously ill hospitalized non-CF patients, signs of nephrotoxicity have been reported in approximately 20% of patients.

- Hypersensitivity reactions including skin rash and drug fever have been reported. If these occur treatment should be withdrawn.
- Local irritation at the site of injection may occur.

#### Inhalation treatment

- Inhalation may induce coughing or bronchospasm.
- Sore throat or mouth has been reported and may be due to *Candida albicans* infection of hypersensitivity. Skin rash may also indicate hypersensitivity, if this occurs treatment should be withdrawn.

#### **4.9 Overdose**

Overdose can result in neuromuscular blockade that can lead to muscular weakness, apnoea and possible respiratory arrest. Overdose can also cause acute renal failure characterized by decreased urine output and increase serum concentrations of BUN and creatinine.

There is no specific antidote, manage by supportive treatment. Measures to increase the rate of elimination of colistin e.g. mannitol diuresis, prolonged haemodialysis or peritoneal dialysis may be tried, but effectiveness is unknown.

### **5. Pharmacological properties**

#### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: antibacterials for systemic use, other antibacterials, polymyxins.

ATC Code: J01XB01

#### Mechanism of action

Colistin is a cyclic polypeptide antibacterial agent belonging to the polymyxin group. Polymyxins work by damaging the cell membrane and the resulting physiological effects are lethal to the bacterium. Polymyxins are selective for aerobic Gram-negative bacteria that have a hydrophobic outer membrane.

#### Resistance

Resistance bacteria are characterized by modification of the phosphate groups of lipopolysaccharide, which become substituted with ethanolamine or aminoarabinose. Naturally resistant Gram-negative bacteria, such as *Proteus mirabilis* and *Burkholderia cepacian*, show complete substitution of their lipid phosphate by ethanolamine or aminoarabinose.

Cross resistance between colistin (polymyxin E) and polymyxin B is expected. Since the mechanism of action of the polymyxins is different from that of other antibacterial agents, resistance to colistin and polymyxin by the above mechanism alone would not be expected to result in resistance to other drug classes.

#### PK/PD relationship

Polymyxins have been reported to have a concentration-dependent bactericidal effect on susceptible bacteria, fAUC/MIC is considered to be correlated with clinical efficacy.

| <b>EUCAST Breakpoints</b> | <b>Susceptible (S)</b> | <b>Resistant (R)<sup>a</sup></b> |
|---------------------------|------------------------|----------------------------------|
| <i>Acinetobacter</i>      | S $\leq$ 2             | R > 2 mg/L                       |
| <i>Enterobacteriaceae</i> | S $\leq$ 2             | R > 2 mg/L                       |
| <i>Pseudomonas</i> spp    | S < 4                  | R > 2 mg/L                       |

<sup>a</sup> Breakpoints apply to dosage of 2-3 MIU (66.66-100 mg) x 3. A loading (9 MIU (300 mg)) may be needed.

### Susceptibility

The prevalence of acquired resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable.

| <b>Commonly susceptible species</b>                  |
|------------------------------------------------------|
| <i>Acinetobacter calcoaceticus-baumannii complex</i> |
| <i>Klebsiella aerogenes / Klebsiella pneumoniae</i>  |
| <i>Escherichia coli</i>                              |
| <i>Pseudomonas aeruginosa</i>                        |
| <b>Inherently resistant organisms</b>                |
| <i>Burkholderia cepacian and related species</i>     |
| <i>Morganella morganii</i>                           |
| <i>Proteus mirabilis / Proteus vulgaris</i>          |
| <i>Providencia rettgeri / Providencia stuartii</i>   |
| <i>Serratia species</i>                              |

## 5.2 Pharmacokinetic properties

### Absorption

The information on the pharmacokinetics of colistimethate sodium (CMS) and colistin is limited. There are indications that pharmacokinetics in critically ill patients differ from those in patients with less severe physiological derangement and from those in healthy volunteers. The following data are based on studies using HPLC to determine CMS/colistin plasma concentrations.

After infusion of colistimethate sodium the inactive pro-drug is converted to the active colistin. Peak plasma concentrations of colistin have been shown to occur with a delay of up to 7 hours after administration of colistimethate sodium in critically ill patients.

Absorption from the gastrointestinal tract does not occur to any appreciable extent in the normal individual.

When given by nebulization, variable absorption has been reported that may depend on the aerosol particle size, nebuliser system and lung status. Studies in healthy volunteers and patients with various infection have reported serum levels from nil to potentially therapeutic concentrations of 4mg/l or more. Therefore, the possibility of systemic absorption should always be borne in mind when treating patients by inhalation.

### Distribution

The volume of distribution of colistin in healthy subjects is low and corresponds approximately to extracellular fluid (ECF). The volume of distribution is relevantly enlarged in critically ill subjects. Protein binding is moderate and decreases at higher concentrations. In the absence of meningeal inflammation, penetration into the cerebrospinal fluid (CSF) is minimal, but increase in the presence of meningeal inflammation.

Both CMS and colistin display linear PK in the clinically relevant dose range.

### Elimination

It is estimated that approximately 30% of colistimethate sodium is converted to colistin in healthy subjects, its clearance is dependent on creatinine clearance and as renal function decreases, a greater portion of CMS is converted to colistin. In patients with very poor renal function (creatinine clearance <30 ml/min), the extent of conversion could be as high as 60 to 70%. CMS is eliminated predominantly by the kidneys via glomerular filtration. In healthy subjects, 60% to 70% of CMS is excreted unchanged in the urine within 24 hours.

The elimination of the active colistin is incompletely characterized. Colistin undergoes extensive renal tubular reabsorption and may either be cleared non-renally or undergo renal metabolism with the potential for renal accumulation. Colistin clearance is decreased in renal impairment, possibly due to increased conversion of CMS.

Half-life of colistin in healthy subjects and those with cystic fibrosis is reported to be around 3h and 4h, respectively, with a total clearance of around 3L/h. In critically ill patients, half-life has been reported to be prolonged to around 9-18h.

## **5.3 Preclinical safety data**

Data on potential genotoxicity are limited and carcinogenicity data for colistimethate sodium are lacking. Colistimethate sodium has been shown to induce chromosomal aberrations in human lymphocytes, *in vitro*. This effect may be related to a reduction in mitotic index, which was also observed.

Reproductive toxicity studies in rats and mice do not indicate teratogenic properties. However, colistimethate sodium given intramuscularly during organogenesis to rabbits at 4.15 and 9.3 mg/kg resulted in talipes varus in 2.6 and 2.9% of fetuses respectively. These doses are 0.5 and 1.2 times the maximum daily human dose. In addition, increased resorption occurred at 9.3 mg/kg.

There are no other preclinical safety data of relevance to the prescriber which are additional to safety data derived from patient exposure and already included in other sections of the SPC.

## 6. Pharmaceutical Particulars

**6.1 List of excipients:** None

**6.2 Incompatibilities:** Mix infusion and nebuliser solution involving colistimethate sodium should be avoided.

**6.3 Shelf life:** Before opening: 2 years

The chemical and physical in-use stability of reconstituted with 2ml sterile water for injection or 0.9% sodium chloride or 5% Dextrose in 0.9% sodium chloride or 5% Dextrose in water in the vial, with a concentration 75 mg/ml, has been demonstrated for 7 days in refrigerator (2-8°C) or store below 30°C.

The chemical and physical in-use stability of dilution with 100ml 0.9% sodium chloride or 5% Dextrose in water or 5% Dextrose in 0.225% sodium chloride or 5% Dextrose in 0.45% sodium chloride or 5% Dextrose in 0.9% sodium chloride or Lactate Ringer's or 10% invert sugar, with a concentration 1.5 mg/ml, has been demonstrated for 24 hours in store below 30°C.

From a microbiological point of view, unless the method of opening/ reconstitution/ dilution precludes the risk of microbial contamination, the product should be used immediately.

If not used immediately, in-use storage times and condition are the responsibility of user.

For solution for intrathecal and intraventricular administration, the reconstituted product should be used immediately.

For solution for inhalation by nebuliser, the reconstituted product should be freshly prepared before use.

### 6.4 Special precautions for storage

Store below 30°C

Keep the vials in the outer carton in order to protect from light.

### 6.5 Nature and contents of container

Clear glass vial (Type I) with grey chlorobutyl rubber stopper, sealed with aluminium/polypropylene flip-off cap contains Colistimethate Sodium equivalent to Colistin 150 mg, packed or unpacked in a box of 1, 2, 3, 4, 5, 10, 12, 20, 24, 25, 50 and 100 vials.

### 6.6 Special precaution for storage

For bolus injection: Reconstitute the contents of the vial with not more than 10ml sterile water for injection or 0.9% sodium chloride.

For infusion: The contents of the reconstituted vial may be diluted, usually with 50ml 0.9% sodium chloride.

When the intrathecal and intraventricular routes of administration are used, the volume administered should not exceed 1 ml (reconstituted concentration 125,000 IU (0.4 mg of CBA)/ml).

For inhalation by nebuliser: Reconstitute the contents of the vial with either water for injections to produce a hypotonic solution or a 50:50 mixture of water for injections and 0.9% sodium chloride to produce an isotonic solution or with 0.9% sodium chloride to produce a hypertonic solution.

The volume of reconstitution should be according to the instructions for use of nebuliser administration device, and is normally not more than 4ml.

The output from the nebuliser may be vented to the open air or a filter may be fitted. Nebulisation should take place in a well ventilated room.

During reconstitution swirl gently to avoid frothing.

Solutions are for single use only, should be freshly prepared before used and any remaining solution should be discarded.

**7. Marketing Authorization Holder**

ABLE MEDICAL COMPANY LIMITED

111 Moo. 9 Nong Son, Chiang Yuen,

Mahasarakham 44160, Thailand

**8. Marketing Authorization Numbers**

1A 15122/63

**9. Date of authorization**

21 July, 2020

**10. Date of revision of the text**

12 June, 2023