

Lasiven

AB-424/2024_04

Composition	:	Each 2 mL contains Furosemide 20 mg
Indication	:	Edema Acute pulmonary edema Congestive heart failure (CHF) and chronic renal failure
Dosage and administration	:	Edema : The initial dose is 20 to 40 mg IM or IV. If needed, another dose may be given in the same manner 2 hours later. The dose may be raised by 20 mg and given no sooner than 2 hours after previous dose, until desired diuretic effect is obtained. This dose should be given once or twice daily. Acute pulmonary edema : The initial dose is 40 mg IV (over 1 to 2 minutes). If response is not satisfactory within 1 hour, increase to 80 mg IV. Congestive heart failure (CHF) and chronic renal failure : The dose is 2 to 2.5 g/day or more. For IV bolus injection, the maximum dose should not exceed 1 g/day given over 30 minutes.
Adverse reaction	:	Fluid and electrolyte imbalance including hyponatremia, hypokalemia and hypochloremic alkalosis, particularly after large doses or prolonged administration.
Contraindication	:	Acute renal failure with anuria, hyponatremia, hypokalemia, impaired renal or hepatic function.
Manufactured by	:	ABLE MEDICAL CO., LTD 111 Moo. 9 Nong Son, Chiang Yuen, Mahasarakham 44160, Thailand