

เอกสารกำกับยาภาษาอังกฤษสำหรับแพทย์

1. NAME OF THE MEDICINAL PRODUCT

SPORAL® CAPSULES

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each capsule, for oral administration, contains 100 mg itraconazole in a pellet formulation, and is available as a blue opaque cap and pink transparent body.

For excipients, see section *6.1 List of excipients*.

3. PHARMACEUTICAL FORM

Capsules, hard

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

SPORAL capsules are indicated for the following conditions:

- Gynecological indications:
 - Treatment of vulvovaginal candidosis.
- Dermatological / mucosal / ophthalmological indications:
 - Treatment of dermatomycosis, including highly keratinized regions as in plantar tinea pedis and palmar tinea manus;
 - Treatment of pityriasis versicolor;
 - Treatment of oral candidosis;
 - Treatment of fungal keratitis.
- Treatment of onychomycosis, caused by dermatophytes and/or yeasts.
- Systemic mycoses, only in the following fungal infections:
 - Treatment of systemic aspergillosis and candidosis;
 - Cryptococcosis (including cryptococcal meningitis):

- Treatment of immunocompromised patients with cryptococcosis and in all patients with cryptococcosis of the central nervous system, only when first line treatment is considered inappropriate or has proven ineffective;
- For maintenance therapy of cryptococcal meningitis in AIDS patients, only when first line treatment is considered inappropriate or has proven ineffective.
- Treatment of histoplasmosis;
- Histoplasmosis, maintenance therapy only in AIDS patients;
- Treatment of blastomycosis;
- Treatment of sporotrichosis, including lymphocutaneous/cutaneous and extracutaneous;
- Treatment of paracoccidioidomycosis;
- Treatment of chromoblastomycosis (formerly chromomycosis).

4.2 Posology and method of administration

For optimal absorption, administer SPORAL capsules immediately after a full meal.

The capsules must be swallowed whole.

Gynecological indication

Indication	Dose	Treatment Duration
Treatment of vulvovaginal candidosis	200 mg twice daily or 200 mg once daily	1 day or 3 days

Dermatological / mucosal / ophthalmological indications		
Indication	Dose	Treatment Duration
Treatment of dermatomycosis	200 mg once daily or 100 mg once daily	7 days or 15 days
Treatment of dermatomycosis in highly keratinized regions as in plantar tinea pedis and palmar tinea manus	200 mg twice daily or 100 mg once daily	7 days or 30 days
Treatment of pityriasis versicolor	100 mg twice daily or 200 mg once daily	5 - 7 days
Treatment of oral candidosis	100 mg once daily	15 days
Treatment of fungal keratitis	200 mg once daily	21 days The duration of treatment should be adjusted to the clinical response.

Onychomycosis, caused by dermatophytes and/or yeasts									
Onychomycosis Pulse treatment		Dose and Treatment duration							
		A pulse treatment consists of two capsules twice daily (200 mg twice daily) for one week. Two pulse treatments are recommended for fingernail infections, and three pulse treatments for toenail infections. Pulse treatments are always separated by a 3-week drug-free interval. Clinical response will become evident as the nail re-grows, following discontinuation of the treatment.							
Site of onychomycosis	Week 1	Week 2	Week 3	Week 4	Week 5	Week 6	Week 7	Week 8	Week 9
Toenails with or without fingernail involvement	Pulse 1	Itraconazole-free weeks			Pulse 2	Itraconazole-free weeks			Pulse 3
Fingernails only	Pulse 1	Itraconazole-free weeks			Pulse 2				
Onychomycosis Continuous treatment		Dose				Treatment duration			
Toenails with or without fingernail involvement		200 mg once daily				3 months			

Elimination of itraconazole from skin and nail tissue is slower than from plasma. Optimal clinical and mycological response is thus reached 2 to 4 weeks after the cessation of treatment for skin infections and 6 to 9 months after the cessation of treatment for nail infections.

Systemic mycoses			
Indication	Dose	Median Treatment Duration¹	Remarks
Treatment of aspergillosis	200 mg once daily	2-5 months	Increase dose to 200 mg twice daily in case of invasive or disseminated disease.
Treatment of candidosis	100 - 200 mg once daily	3 weeks - 7 months	Increase dose to 200 mg twice daily in case of invasive or disseminated disease.
Treatment of non-meningeal cryptococcosis	200 mg once daily	2 months - 1 year	
Treatment of cryptococcal meningitis	200 mg twice daily	2 months -1 year	
Cryptococcal meningitis (maintenance therapy only in AIDS patients)	200 mg once daily	<i>Until immune recovery²</i>	
Treatment of histoplasmosis	200 mg once daily - 200 mg twice daily	8 months	
Histoplasmosis (maintenance therapy only in AIDS patients)	200 mg once or twice daily	<i>Until immune recovery²</i>	
Treatment of blastomycosis	100 mg once daily - 200 mg twice daily	6 months	
Treatment of lymphocutaneous and cutaneous sporotrichosis	100 mg or 200 mg once daily (localized lesions) or 200 mg twice daily (extensive lesions)	3 months to 6 months	
Treatment of extracutaneous sporotrichosis	200 mg twice daily	12 months	
Treatment of paracoccidioidomycosis	100 mg once daily	6 months	Data on the efficacy of SPORAL capsules at this dosage for treatment of paracoccidioidomycosis in patients with AIDS is not available.
Treatment of chromoblastomycosis (formerly chromomycosis)	200 - 400 mg once daily	6 months	

¹ The duration of treatment should be adjusted depending on the clinical response. ² The duration of treatment should be based upon the status of the immune recovery.

Special populations

Pediatrics

Clinical data on the use of SPORAL capsules in pediatric patients are limited. The use of SPORAL capsules in pediatric patients is not recommended unless it is determined that the potential benefit outweighs the potential risks (See section 4.4 *Special warning and precautions for use*).

Elderly

Clinical data on the use of SPORAL capsules in elderly patients are limited. It is advised to use SPORAL capsules in these patients only if it is determined that the potential benefit outweighs the potential risks. In general, it is recommended that the dose selection for an elderly patient should be taken into consideration, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (See section 4.4 *Special warning and precautions for use*).

Hepatic impairment

Limited data are available on the use of oral itraconazole in patients with hepatic impairment. Caution should be exercised when this drug is administered in this patient population (See section 5.2 *Pharmacokinetic Properties - Special Populations, Hepatic impairment*).

Renal impairment

Limited data are available on the use of oral itraconazole in patients with renal impairment. The exposure of itraconazole may be lower in some patients with renal insufficiency. Caution should be exercised when this drug is administered in this patient population and adjusting the dose may be considered.

4.3 Contraindications

- SPORAL capsules are contraindicated in patients with known hypersensitivity to itraconazole or to any of the excipients.
- Coadministration of a number of CYP3A4 substrates is contraindicated with SPORAL capsules. Increased plasma concentrations of these drugs, caused by coadministration with itraconazole, may increase or prolong both therapeutic and adverse effects to such an extent that a potentially serious situation may occur. For example, increased plasma concentrations of some of these drugs can lead to QT prolongation and ventricular tachyarrhythmias including occurrences of torsade de pointes, a potentially fatal arrhythmia. Some specific examples are listed in 4.5 *Interaction with other medicinal products and other forms of interactions*.
- SPORAL capsules should not be administered to patients with evidence of ventricular dysfunction such as congestive heart failure (CHF) or a history of CHF except for the

treatment of life-threatening or other serious infections (See section 4.4 *Special warning and precautions for use*).

- SPORAL capsules must not be used during pregnancy (except for life-threatening cases) (See section 4.6 *Fertility, pregnancy and lactation*).

Women of childbearing potential taking SPORAL should use contraceptive precautions. Highly effective contraception should be continued until the menstrual period following the end of SPORAL therapy.

4.4 Special warning and precautions for use

Cross-hypersensitivity

There is limited information regarding cross-hypersensitivity between itraconazole and other azole antifungal agents. Caution should be used in prescribing SPORAL capsules to patients with hypersensitivity to other azoles.

Cardiac effects

In a healthy volunteer study with SPORAL IV, a transient asymptomatic decrease of the left ventricular ejection fraction was observed; this resolved before the next infusion. The clinical relevance of these findings to the oral formulations is unknown.

Itraconazole has been shown to have a negative inotropic effect and SPORAL has been associated with reports of congestive heart failure. Heart failure was more frequently reported among spontaneous reports of 400 mg total daily dose than among those of lower total daily doses, suggesting that the risk of heart failure might increase with the total daily dose of itraconazole.

SPORAL should not be used in patients with congestive heart failure or with a history of congestive heart failure unless the benefit clearly outweighs the risk. This individual benefit/risk assessment should take into consideration factors such as the severity of the indication, the dosing regimen (e.g., total daily dose), and individual risk factors for congestive heart failure. These risk factors include cardiac disease, such as ischemic and valvular disease; significant pulmonary disease, such as chronic obstructive pulmonary disease; and renal failure and other edematous disorders. Such patients should be informed of the signs and symptoms of congestive heart failure, should be treated with caution, and should be monitored for signs and symptoms of congestive heart failure during treatment; if such signs or symptoms do occur during treatment, SPORAL should be discontinued.

Calcium channel blockers can have negative inotropic effects which may be additive to those of itraconazole. In addition, itraconazole can inhibit the metabolism of calcium channel blockers. Therefore, caution should be used when co-administering itraconazole and calcium channel blockers due to an increased risk of CHF.

Hepatic effects

Very rare cases of serious hepatotoxicity, including some cases of fatal acute liver failure, have occurred with the use of SPORAL. Most of these cases involved patients who, had

pre-existing liver disease, were treated for systemic indications, had significant other medical conditions and/or were taking other hepatotoxic drugs. Some patients had no obvious risk factors for liver disease. Some of these cases were observed within the first month of treatment, including some within the first week. Liver function monitoring should be considered in patients receiving SPORAL treatment. Patients should be instructed to promptly report to their physician signs and symptoms suggestive of hepatitis such as anorexia, nausea, vomiting, fatigue, abdominal pain or dark urine. In these patients treatment should be stopped immediately and liver function testing should be conducted.

Limited data are available on the use of oral itraconazole in patients with hepatic impairment. Caution should be exercised when the drug is administered in this patient population. It is recommended that patients with impaired hepatic function be carefully monitored when taking itraconazole. It is recommended that the prolonged elimination half-life of itraconazole observed in the single oral dose clinical trial with itraconazole capsules in cirrhotic patients be considered when deciding to initiate therapy with other medications metabolized by CYP3A4.

In patients with elevated or abnormal liver enzymes or active liver disease, or who have experienced liver toxicity with other drugs, treatment with SPORAL is strongly discouraged unless there is a serious or life-threatening situation where the expected benefit exceeds the risk. It is recommended that liver function monitoring be done in patients with pre-existing hepatic function abnormalities or those who have experienced liver toxicity with other medications.

(See section *5.2 Pharmacokinetic Properties - Special Populations, Hepatic impairment.*)

Reduced gastric acidity

Absorption of itraconazole from SPORAL capsules is impaired when gastric acidity is reduced. In patients with reduced gastric acidity, whether from disease (e.g. patients with achlorhydria) or from concomitant medication (e.g. patients taking drugs that reduce gastric acidity), it is advisable to administer SPORAL capsules with an acidic beverage (such as non-diet cola). The antifungal activity should be monitored and the itraconazole dose increased as deemed necessary (See section *4.5 Interaction with other medicinal products and other forms of interactions* and section *5.2 Pharmacokinetic properties - Absorption*).

Pediatrics

Clinical data on the use of SPORAL capsules in pediatric patients are limited. The use of SPORAL capsules in pediatric patients is not recommended unless it is determined that the potential benefit outweighs the potential risks.

Elderly

Clinical data on the use of SPORAL capsules in elderly patients are limited. It is advised to use SPORAL capsules in these patients only if it is determined that the potential benefit outweighs the potential risks. In general, it is recommended that the dose selection for an elderly patient should be taken into consideration, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Renal impairment

Limited data are available on the use of oral itraconazole in patients with renal impairment. The exposure of itraconazole may be lower in some patients with renal insufficiency. Caution should be exercised when this drug is administered in this patient population and adjusting the dose may be considered.

Hearing loss

Transient or permanent hearing loss has been reported in patients receiving treatment with itraconazole. Several of these reports included concurrent administration of quinidine which is contraindicated (See section *4.3 Contraindications*). The hearing loss usually resolves when treatment is stopped, but can persist in some patients.

Immunocompromised patients

In some immunocompromised patients (e.g., neutropenic, AIDS or organ transplant patients), the oral bioavailability of SPORAL capsules may be decreased. Therefore, the dose should be adjusted based on the clinical response in these patients.

Patients with immediately life-threatening systemic fungal infections

Due to the pharmacokinetic properties (See section *5.2 Pharmacokinetic Properties*), SPORAL capsules are not recommended for initiation of treatment in patients with immediately life-threatening systemic fungal infections.

Patients with AIDS

In patients with AIDS who have received treatment for a systemic fungal infection with SPORAL capsules and who are considered at risk for relapse, the treating physician should evaluate the need for a maintenance treatment.

Neuropathy

If neuropathy occurs that may be attributable to SPORAL capsules, the treatment should be discontinued.

Disorders of Carbohydrate Metabolism

Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

Cross-resistance

In systemic candidosis, if fluconazole-resistant strains of *Candida* species are suspected, it cannot be assumed that these are sensitive to itraconazole, hence it is recommended to have their sensitivity tested before the start of itraconazole therapy.

Interaction potential

Coadministration of specific drugs with itraconazole may result in changes in efficacy of itraconazole and/or the coadministered drug, life-threatening effects and/or sudden death. Drugs that are contraindicated, not recommended or recommended for use with caution in combination with itraconazole are listed in *4.5 Interaction with other medicinal products and other forms of interactions*.

Cystic fibrosis

In cystic fibrosis patients, variability in therapeutic levels of itraconazole was observed with steady state dosing of itraconazole oral solution using 2.5 mg/kg bid. Steady state concentrations of > 250 ng/mL were achieved in approximately 50% of subjects greater than 16 years of age, but in none of the patients less than 16 years of age. If a patient does not respond to SPORAL capsules, consideration should be given to switching to alternative therapy.

4.5 Interaction with other medicinal products and other forms of interactions

Itraconazole is a drug with a high interaction potential. The various types of interaction and associated general recommendations are described below. In addition, a table is provided listing examples of drugs that may interact with itraconazole, organized per drug family for easy reference.

Itraconazole is mainly metabolized through CYP3A4. Other substances that either share this metabolic pathway or modify CYP3A4 activity may influence the pharmacokinetics of itraconazole. Coadministration of itraconazole with moderate or potent CYP3A4 inducers may decrease the bioavailability of itraconazole and hydroxy-itraconazole to such an extent that efficacy may be reduced. Coadministration with moderate or potent inhibitors of CYP3A4 may increase the bioavailability of itraconazole, which may result in increased or prolonged pharmacologic effects of itraconazole.

Absorption of itraconazole from the capsule formulation is reduced in subjects with reduced gastric acidity. Drugs that reduce gastric acidity impair the absorption of itraconazole from itraconazole capsules. To counteract this effect, it is recommended to administer itraconazole capsules with an acidic beverage (such as non-diet cola) upon coadministration with drugs that reduce gastric acidity (see section *4.4 Special warning and precautions for use*).

Itraconazole and its major metabolite, hydroxy-itraconazole are potent CYP3A4 inhibitors. Itraconazole is an inhibitor of the drug transporters P-glycoprotein and breast cancer resistance protein (BCRP). Itraconazole can inhibit the metabolism of drugs metabolized by CYP3A4 and can inhibit the drug transport by P-glycoprotein and/or BCRP, which may result in increased plasma concentrations of these drugs and/or their active metabolite(s) when they are administered with itraconazole. These elevated plasma concentrations may increase or prolong both therapeutic and adverse effects of these drugs. For some drugs, coadministration with itraconazole may result in decreased plasma concentrations of the drug or of the active moiety of the drug. This may result in reduced efficacy of the drug.

Following cessation of medical treatment with itraconazole, plasma concentrations decrease below the detection limit within 7 to 14 days, depending on the dose and duration of treatment. In patients with hepatic cirrhosis or in subjects receiving CYP3A4 inhibitors the plasma concentrations decline slower. This is particularly important for consideration when initiating therapy with drugs whose metabolism is affected by itraconazole.

The following general recommendations apply, unless stated differently in table.

- 'Contraindicated': Under no circumstances is the drug to be coadministered with itraconazole. This applies to:
 - CYP3A4 substrates for which increased plasma concentrations may increase or prolong therapeutic and/or adverse effects to such an extent that a potentially serious situation may occur (see section 4.3 *Contraindications*).
- 'Not recommended': It is recommended that the use of the drug be avoided, unless the benefits outweigh the potentially increased risks. If coadministration cannot be avoided, clinical monitoring is recommended, and the dosage of itraconazole and/or the coadministered drug adapted as deemed necessary. When appropriate, it is recommended that plasma concentrations be measured. This applies to:
 - Moderate or potent CYP3A4 inducers: not recommended from 2 weeks before and during treatment with itraconazole
 - CYP3A4/P-gp/BCRP substrates for which increased or decreased plasma concentrations result in significant risk: not recommended during and up to 2 weeks after treatment with itraconazole.
- 'Use with caution': Careful monitoring is recommended when the drug is coadministered with itraconazole. Upon coadministration, it is recommended that patients be monitored closely and the dosage of itraconazole and/or the coadministered drug adapted as deemed necessary. When appropriate, it is recommended that plasma concentrations be measured. This applies to:
 - Drugs that reduce gastric acidity (itraconazole capsules only)
 - Moderate or potent inhibitors of CYP3A4
 - CYP3A4/P-gp/BCRP substrates for which increased or decreased plasma concentrations result in a clinically relevant risk

The list of examples of interacting drugs in the table below is not comprehensive and therefore the label of each drug that is coadministered with itraconazole should be consulted for information related to the route of metabolism, interaction pathways, potential risks, and specific actions to be taken with regards to coadministration. The drugs listed in this table are based on either drug interaction studies or case reports, or potential interactions based on the mechanism of interaction.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
Alpha Blockers		
Alfuzosin Silodosin Tamsulosin	Alfuzosin C _{max} (↑↑), AUC (↑↑) ^a Silodosin C _{max} (↑↑), AUC (↑↑) ^a Tamsulosin C _{max} (↑↑), AUC (↑↑) ^a	Not recommended during and for 2 weeks after treatment with itraconazole. Increased risk of alfuzosin/silodosin/tamsulosin-related adverse reactions ^c .
Analgesics		
Alfentanil Buprenorphine (IV and sublingual) Oxycodone Sufentanil	Alfentanil AUC (↑↑ to ↑↑↑↑) ^a Buprenorphine C _{max} (↑↑), AUC (↑↑) ^a Oxycodone C _{max} ↑, AUC ↑↑ Sufentanil conc increase (extent unknown) ^{a,b}	Use with caution, monitor for adverse reactions related to the analgesic ^c , dose reduction of alfentanil/buprenorphine/oxycodone/sufentanil may be necessary.
Fentanyl	Fentanyl IV AUC (↑↑) ^a Fentanyl other form. conc increase (extent unknown) ^{a,b}	Not recommended during and for 2 weeks after treatment with itraconazole. Increased risk of fentanyl-related adverse reactions ^c .
Levacetylmethadol (levomethadyl)	Levacetylmethadol C _{max} (↑↑), AUC (↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of levacetylmethadol-related adverse reactions, such as QT prolongation and TdP.
Methadone	(R)-methadone C _{max} (↑), AUC (↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of methadone-related adverse reactions, such as potentially life-threatening respiratory depression, QT prolongation and TdP.
Antiarrhythmics		
Digoxin	Digoxin C _{max} ↑, AUC ↑	Use with caution, monitor for digoxin adverse reactions, dose reduction of digoxin may be necessary ^c .
Disopyramide	Disopyramide conc increase (↑↑) ^{a,b}	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of disopyramide-related

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
		adverse reactions, such as serious arrhythmias including TdP.
Dofetilide	Dofetilide C_{max} (\uparrow), AUC (\uparrow) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of dofetilide-related adverse reactions, such as serious ventricular arrhythmias including TdP.
Dronedarone	Dronedarone C_{max} ($\uparrow\uparrow\uparrow$), AUC ($\uparrow\uparrow\uparrow$) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of dronedarone-related adverse reactions, such as QT prolongation and cardiovascular death.
Quinidine	Quinidine C_{max} \uparrow , AUC $\uparrow\uparrow$	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of quinidine-related adverse reactions, such as QT prolongation, TdP, hypotension, confusion and delirium.
Antibacterials		
Bedaquiline	Bedaquiline C_{max} (\leftrightarrow), AUC (\uparrow) during 2 weeks of bedaquiline q.d. dosing ^a	Not recommended, coadministration for more than 2 weeks at any time during bedaquiline dosing is not recommended: increased risk of bedaquiline-related adverse reactions ^c
Ciprofloxacin Erythromycin	Itraconazole C_{max} \uparrow , AUC \uparrow	Use with caution, monitor for itraconazole adverse reactions, dose reduction of itraconazole may be necessary.
Clarithromycin	Clarithromycin conc increase (extent unknown) ^{a,b} Itraconazole C_{max} \uparrow , AUC \uparrow ;	Use with caution, monitor for adverse reactions related to itraconazole and/or clarithromycin ^c , dose reduction of itraconazole and/or clarithromycin may be necessary.
Delamanid Trimetrexate	Delamanid conc. increase (extent unknown) ^{a,b} Trimetrexate conc increase (extent unknown) ^{a,b}	Use with caution, monitor for delamanid/trimetrexate adverse reactions ^c , dose reduction of delamanid/trimetrexate may be necessary
Isoniazid Rifampicin	Isoniazid: itraconazole conc. ($\downarrow\downarrow\downarrow$) ^{a,b}	Not recommended from 2 weeks before and during treatment with itraconazole, Itraconazole efficacy may be reduced.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
	Rifampicin: itraconazole AUC ↓↓↓	
Rifabutin	Rifabutin conc. increase (extent unknown) ^{a,b} Itraconazole: C _{max} ↓↓, AUC ↓↓	Not recommended from 2 weeks before, during and for 2 weeks after treatment with itraconazole. Itraconazole efficacy may be reduced and increased risk of rifabutin-related adverse reactions ^c .
Telithromycin	In healthy subjects: telithromycin C _{max} ↑, AUC ↑ In severe renal impairment: telithromycin AUC (↑↑) ^a In severe hepatic impairment: telithromycin conc. increase (extent unknown) ^{a,b}	Contraindicated in patients with severe renal or hepatic impairment during and for 2 weeks after treatment with itraconazole, Increased risk of telithromycin-related adverse reactions ^c , such as hepatotoxicity, QT prolongation and TdPs. Use with caution in other patients; monitor for telithromycin adverse reactions, dose reduction of telithromycin may be necessary
Anticoagulants and Antiplatelet Drugs		
Apixaban Edoxaban Rivaroxaban Vorapaxar	Apixaban C _{max} (↑), AUC (↑) ^a Edoxaban C _{max} (↑), AUC (↑) ^a Rivaroxaban C _{max} (↑), AUC (↑ to ↑↑) ^a Vorapaxar C _{max} (↑), AUC (↑) ^a	Not recommended during and for 2 weeks after treatment with itraconazole. Increased risk of apixaban/edoxaban/rivaroxaban/vorapaxar-related adverse reactions ^c .
Coumarins (eg, warfarin) Cilostazol	Coumarins (eg, warfarin) conc increase (extent unknown) ^{a,b} Cilostazol C _{max} (↑), AUC (↑↑) ^a	Use with caution, monitor for coumarins/cilostazol adverse reactions ^c , dose reduction of coumarins/cilostazol may be necessary.
Dabigatran	Dabigatran C _{max} (↑↑), AUC (↑↑) ^a	Use with caution, monitor for dabigatran adverse reactions ^c , dose reduction of dabigatran may be necessary
Ticagrelor	Ticagrelor C _{max} (↑↑), AUC (↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
		Increased risk of ticagrelor-related adverse reactions, such as bleeding.
Anticonvulsants		
Carbamazepine	Carbamazepine conc. (↑) ^{a,b} Itraconazole conc. (↓↓) ^{a,b}	Not recommended from 2 weeks before, during and for 2 weeks after treatment with itraconazole. Itraconazole efficacy may be reduced and increased risk for carbamazepine-related adverse reactions ^c .
Phenobarbital Phenytoin	Phenobarbital: itraconazole conc. (↓↓) ^{a,b} Phenytoin: itraconazole AUC ↓↓↓	Not recommended from 2 weeks before and during treatment with itraconazole. Itraconazole efficacy may be reduced.
Antidiabetics		
Repaglinide Saxagliptin	Repaglinide C _{max} ↑, AUC ↑ Saxagliptin C _{max} (↑↑), AUC (↑↑) ^a	Use with caution, monitor for repaglinide/saxagliptin adverse reactions ^c , dose reduction of repaglinide/saxagliptin may be necessary
Antihelminthics, antifungals and antiprotozoals		
Artemether-lumefantrine Quinine	Artemether C _{max} (↑↑), AUC (↑↑) ^a Lumefantrine C _{max} (↑), AUC (↑) ^a Quinine C _{max} ↔, AUC ↑	Use with caution, monitor for artemether-lumefantrine/quinine adverse reactions ^c . Refer to the label for specific actions to be taken.
Halofantrine	Halofantrine conc increase (extent unknown) ^{a,b}	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of halofantrine-related adverse reactions, such as QT prolongation and fatal arrhythmias.
Isavuconazole	Isavuconazole C _{max} (↔), AUC (↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of isavuconazole-related adverse reactions, such as hepatic adverse reactions, hypersensitivity reactions and embryo-fetal toxicity.
Praziquantel	Praziquantel C _{max} (↑↑), AUC (↑) ^a	Use with caution, monitor for praziquantel adverse reactions ^c , dose reduction of praziquantel may be necessary.
Antihistamines		
Astemizole	Astemizole C _{max} (↑), AUC (↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
		Increased risk of astemizole-related adverse reactions, such as QT prolongation, TdP and other ventricular arrhythmias.
Bilastine Ebastine Rupatadine	Bilastine C _{max} (↑↑), AUC (↑) ^a Ebastine C _{max} ↑↑, AUC ↑↑↑ Rupatadine conc increase (↑↑↑↑) ^{a,b}	Use with caution, monitor for bilastine/ebastine/rupatadine adverse reactions ^c , dose reduction of bilastine/ebastine/rupatadine may be necessary.
Mizolastine	Mizolastine C _{max} (↑), AUC (↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of mizolastine-related adverse reactions, such as QT prolongation.
Terfenadine	Terfenadine conc increase (extent unknown) ^b	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of terfenadine-related adverse reactions, such as QT prolongation, TdP and other ventricular arrhythmias.
Antimigraine Drugs		
Eletriptan	Eletriptan C _{max} (↑↑), AUC (↑↑↑) ^a	Use with caution, monitor for eletriptan adverse reactions ^c , dose reduction of eletriptan may be necessary.
Ergot alkaloids (such as dihydroergotamine, ergometrine, ergotamine, methylergometrine)	Ergot alkaloids conc increase (extent unknown) ^{a,b}	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of ergot alkaloid-related adverse reactions, such as ergotism.
Antineoplastics		
Bortezomib Brentuximab vedotin Busulfan Erlotinib Gefitinib Imatinib Ixabepilone Nintedanib Panobinostat Pemigatinib Ponatinib Ruxolitinib Sonidegib Tretinoin (oral) Vandetanib	Bortezomib AUC (↑) ^a Brentuximab vedotin AUC (↑) ^a Busulfan C _{max} ↑, AUC ↑ Erlotinib C _{max} (↑↑), AUC (↑) ^a Gefitinib C _{max} ↑, AUC ↑ Imatinib C _{max} (↑), AUC (↑) ^a Ixabepilone C _{max} (↔), AUC (↑) ^a	Use with caution, monitor for adverse reactions related to the antineoplastic drug ^c , dose reduction of the antineoplastic drug may be necessary.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
	Nintedanib C_{max} (↑), AUC (↑) ^a Panobinostat C_{max} (↑), AUC (↑) ^a Pemigatinib C_{max} ↑, AUC ↑ Ponatinib C_{max} (↑), AUC (↑) ^a Ruxolitinib C_{max} (↑), AUC (↑) ^a Sonidegib C_{max} (↑), AUC (↑↑) ^a Tretinoin C_{max} (↑), AUC (↑) ^a Vandetanib C_{max} ↔, AUC ↑	
Idelalisib	Idelalisib C_{max} (↑), AUC (↑) ^a Itraconazole serum conc. increase (extent unknown) ^{a,b}	Use with caution, monitor for adverse reactions related to itraconazole and/or idelalisib ^c , dose reduction of itraconazole and/or idelalisib may be necessary.
Axitinib Bosutinib Cabazitaxel Cabozantinib Ceritinib Cobimetinib Crizotinib Dabrafenib Dasatinib Docetaxel Entrectinib Glasdegib Ibrutinib Lapatinib Nilotinib Olaparib Pazopanib Sunitinib Talazoparib Trabectedin Trastuzumab emtansine Vinca alkaloids	Axitinib C_{max} (↑), AUC (↑↑) ^a Bosutinib C_{max} (↑↑↑), AUC (↑↑↑) ^a Cabazitaxel C_{max} (↔), AUC (↔) ^a Cabozantinib C_{max} (↔), AUC (↑) ^a Ceritinib C_{max} (↑), AUC (↑↑) ^a Cobimetinib C_{max} ↑↑, AUC ↑↑↑ Crizotinib C_{max} (↑), AUC (↑↑) ^a Dabrafenib AUC (↑) ^a Dasatinib C_{max} (↑↑), AUC (↑↑) ^a Docetaxel AUC (↔ to ↑↑) ^a Entrectinib C_{max} ↑, AUC ↑↑↑ Glasdegib C_{max} (↑), AUC (↑↑) ^a	Not recommended during and for 2 weeks after treatment with itraconazole. Increased risk of adverse reactions related to the antineoplastic drug ^c . Additionally: For cabazitaxel, even though the change in pharmacokinetic parameters did not reach statistical significance in a low-dose drug interaction study with ketoconazole, a high variability in the results was observed. For ibrutinib, refer to the label for specific actions to be taken.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
	Ibrutinib C_{max} (↑↑↑↑), AUC (↑↑↑↑) ^a Lapatinib C_{max} (↑↑), AUC (↑↑) ^a Nilotinib C_{max} (↑), AUC (↑↑) ^a Olaparib C_{max} ↑, AUC ↑↑ Pazopanib C_{max} (↑), AUC (↑) ^a Sunitinib C_{max} (↑), AUC (↑) ^a Talazoparib C_{max} ↑, AUC ↑ Trabectedin C_{max} (↑), AUC (↑) ^a Trastuzumab emtansine conc increase (extent unknown) ^{a,b} Vinca alkaloid conc increase (extent unknown) ^{a,b}	
Regorafenib	Regorafenib AUC (↓↓ by estimation of active moiety) ^a	Not recommended during and for 2 weeks after treatment with itraconazole. Regorafenib efficacy may be reduced.
Irinotecan	Irinotecan and its active metabolite conc increase (extent unknown) ^{a,b}	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of irinotecan-related adverse reactions, such as potentially life-threatening myelosuppression and diarrhea.
Mobocertinib	Mobocertinib C_{max} ↑↑, AUC ↑↑↑	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of mobocertinib-related adverse reactions ^c .
Venetoclax	Venetoclax C_{max} (↑↑↑), AUC (↑↑↑) ^a	Contraindicated for chronic lymphocytic leukemia/small lymphocytic lymphoma patients during dose initiation/titration/ramp-up phase of venetoclax. Otherwise, not

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
		recommended during and for 2 weeks after treatment with itraconazole. ^c
Antipsychotics, Anxiolytics and Hypnotics		
Alprazolam Aripiprazole Brotizolam Buspirone Cariprazine Haloperidol Midazolam (iv) Perospirone Quetiapine Ramelteon Risperidone Suvorexant Zopiclone	Alprazolam C _{max} ↔, AUC ↑↑ Aripiprazole C _{max} ↑, AUC ↑ Brotizolam C _{max} ↔, AUC ↑↑ Buspirone C _{max} ↑↑↑↑, AUC ↑↑↑↑ Cariprazine (↑↑) ^{a,b} Haloperidol C _{max} ↑, AUC ↑ Midazolam (iv) conc increase ↑↑ ^b Perospirone C _{max} ↑↑↑, AUC ↑↑↑ Quetiapine C _{max} (↑↑), AUC (↑↑↑) ^a Ramelteon C _{max} (↑), AUC (↑) ^a Risperidone conc increase ↑ ^b Suvorexant C _{max} (↑), AUC (↑↑) ^a Zopiclone C _{max} ↑, AUC ↑	Use with caution, monitor for adverse reactions related to the antipsychotic, anxiolytic or hypnotic drug ^c , dose reduction of these drugs may be necessary.
Lurasidone	Lurasidone C _{max} (↑↑↑), AUC (↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of lurasidone-related adverse reactions, such as hypotension, circulatory collapse, severe extrapyramidal symptoms, seizures.
Midazolam (oral)	Midazolam (oral) C _{max} ↑ to ↑↑, AUC ↑↑ to ↑↑↑↑	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of midazolam-related adverse reactions, such as respiratory depression, cardiac arrest, prolonged sedation and coma.
Pimozide	Pimozide C _{max} (↑), AUC (↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of pimozide-related adverse reactions, such as cardiac

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
		arrhythmias, possibly associated with QT prolongation and TdP.
Sertindole	Sertindole conc increase (extent unknown) ^{a,b}	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of sertindole-related adverse reactions, such as QT prolongation and TdP.
Triazolam	Triazolam C _{max} ↑ to ↑↑, AUC ↑↑ to ↑↑↑	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of triazolam-related adverse reactions, such as seizures, respiratory depression, angioedema, apnea and coma.
Antivirals		
Asunaprevir (boosted) Tenofovir disoproxil fumarate (TDF)	Asunaprevir C _{max} (↑↑↑), AUC (↑↑↑) ^a Tenofovir conc increase (extent unknown) ^{a,b}	Use with caution, however, refer to the label of the antiviral drug for specific actions to be taken.
Boceprevir	Boceprevir C _{max} (↑), AUC (↑↑) ^a Itraconazole conc increase (extent unknown) ^{a,b}	Use with caution, monitor for adverse reactions related to itraconazole and/or boceprevir ^c , dose reduction of itraconazole may be necessary. Refer to the boceprevir label for specific actions to be taken.
Cobicistat	Cobicistat conc increase (extent unknown) ^{a,b} Itraconazole conc increase (extent unknown) ^{a,b}	Use with caution, monitor for adverse reactions related to itraconazole, dose reduction of itraconazole may be necessary.
Daclatasvir Vaniprevir	Daclatasvir C _{max} (↑), AUC (↑↑) ^a Vaniprevir C _{max} (↑↑↑), AUC (↑↑↑) ^a	Use with caution, monitor for daclatasvir/vaniprevir adverse reactions ^c , dose reduction of daclatasvir/vaniprevir may be necessary.
Darunavir (boosted) Fosamprenavir (ritonavir-boosted)	Ritonavir-boosted darunavir: itraconazole C _{max} (↑↑), AUC (↑↑) ^a Ritonavir-boosted fosamprenavir: itraconazole C _{max} (↑), AUC (↑↑) ^a	Use with caution, monitor for itraconazole adverse reactions, dose reduction of itraconazole may be necessary.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
Elvitegravir (boosted)	Elvitegravir C_{max} (\uparrow), AUC (\uparrow) ^a Itraconazole conc increase (extent unknown) ^{a,b}	Use with caution, monitor for adverse reactions related to itraconazole and/or elvitegravir (ritonavir-boosted) ^c . Dose reduction of itraconazole may be necessary; refer to the elvitegravir label for specific actions to be taken.
Efavirenz Nevirapine	Efavirenz: itraconazole C_{max} \downarrow , AUC \downarrow Nevirapine: itraconazole C_{max} \downarrow , AUC $\downarrow\downarrow$	Not recommended from 2 weeks before and during treatment with itraconazole. Itraconazole efficacy may be reduced.
Elbasvir/Grazoprevir	Elbasvir C_{max} (\leftrightarrow), AUC (\uparrow) ^a Grazoprevir C_{max} (\leftrightarrow), AUC ($\uparrow\uparrow$) ^a	Use with caution, monitor for adverse reactions related to the co-administered drugs ^c . Refer to the elbasvir/grazoprevir label for specific actions to be taken.
Glecaprevir/Pibrentasvir	Glecaprevir C_{max} ($\uparrow\uparrow$), AUC ($\uparrow\uparrow$ to $\uparrow\uparrow\uparrow$) ^a Pibrentasvir C_{max} (\leftrightarrow to \uparrow), AUC (\leftrightarrow to $\uparrow\uparrow$) ^a	Use with caution, monitor for adverse reactions related to the co-administered drugs ^c . Refer to the glecaprevir/pibrentasvir label for specific actions to be taken.
Indinavir	Itraconazole conc. \uparrow ^b Indinavir C_{max} \leftrightarrow , AUC \uparrow	Use with caution, monitor for adverse reactions related to itraconazole and/or indinavir ^c , dose reduction of itraconazole and/or indinavir may be necessary.
Maraviroc	Maraviroc C_{max} ($\uparrow\uparrow$), AUC ($\uparrow\uparrow\uparrow$) ^a	Use with caution monitor for adverse reactions ^c . Dose reduction of maraviroc may be necessary.
Ombitasvir/Paritaprevir/Ritonavir with or without Dasabuvir	Itraconazole C_{max} (\uparrow), AUC ($\uparrow\uparrow$) ^a Ombitasvir C_{max} (\leftrightarrow), AUC (\uparrow) ^a Paritaprevir C_{max} (\uparrow), AUC ($\uparrow\uparrow$) ^a Ritonavir C_{max} (\uparrow), AUC (\uparrow) ^a Dasabuvir C_{max} (\uparrow), AUC (\uparrow) ^a	Use with caution, monitor for adverse reactions related to itraconazole and/or the antivirals ^c , dose reduction of itraconazole may be necessary. Refer to the label(s) of the coadministered drugs for specific actions to be taken.
Ritonavir	Itraconazole C_{max} (\uparrow), AUC ($\uparrow\uparrow$) ^a Ritonavir C_{max} (\leftrightarrow), AUC (\uparrow) ^a	Use with caution, monitor for adverse reactions related to itraconazole and/or ritonavir ^c , Dose reduction of itraconazole may be necessary; refer

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
		to the ritonavir label for specific actions to be taken.
Saquinavir	Saquinavir (unboosted) C _{max} ↑↑, AUC ↑↑↑ Itraconazole (with boosted saquinavir) C _{max} (↑), AUC (↑↑) ^a	Use with caution, monitor for adverse reactions related to itraconazole and/or saquinavir ^c , Dose reduction of itraconazole may be necessary; refer to the saquinavir label for specific actions to be taken.
Beta Blockers		
Nadolol	Nadolol C _{max} ↑↑, AUC ↑↑	Use with caution, monitor for nadolol adverse reactions ^c . Dose reduction of nadolol may be necessary.
Calcium Channel Blockers		
Bepridil	Bepridil conc increase (extent unknown) ^{a,b}	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of bepridil-related adverse reactions, such as new arrhythmias and TdP type ventricular tachycardia.
Diltiazem	Diltiazem & Itraconazole conc increase (extent unknown) ^{a,b}	Use with caution, monitor for adverse reactions related to itraconazole and/or diltiazem ^c , dose reduction of itraconazole and/or diltiazem may be necessary.
Felodipine Lercanidipine Nisoldipine	Felodipine C _{max} ↑↑↑, AUC ↑↑↑ Lercanidipine AUC (↑↑↑↑) ^a Nisoldipine C _{max} (↑↑↑↑), AUC (↑↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of dihydropyridine-related adverse reactions, such as hypotension and peripheral edema.
Other dihydropyridines Verapamil	Dihydropyridine conc increase (extent unknown) ^{a,b} Verapamil conc increase (extent unknown) ^{a,b}	Use with caution, monitor for dihydropyridine/verapamil adverse reactions ^c , dose reduction of dihydropyridine/verapamil may be necessary.
Cardiovascular Drugs, Misc		
Aliskiren Riociguat Sildenafil (pulmonary hypertension)	Aliskiren C _{max} ↑↑↑, AUC ↑↑↑ Riociguat C _{max} (↑), AUC (↑↑) ^a	Not recommended during and for 2 weeks after treatment with itraconazole ^c . Increased risk of adverse reactions related to the cardiovascular drug.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
Tadalafil (pulmonary hypertension)	Sildenafil/Tadalafil conc increase (extent unknown but effect may be greater than reported under Urological Drugs) ^{a,b}	
Bosentan Guanfacine	Bosentan C _{max} (↑↑), AUC (↑↑) ^a Guanfacine C _{max} (↑), AUC (↑↑) ^a	Use with caution, monitor for bosentan/guanfacine adverse reactions ^c , dose reduction of bosentan/guanfacine may be necessary.
Ivabradine	Ivabradine C _{max} (↑↑), AUC (↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of ivabradine-related adverse reactions, such as atrial fibrillation, bradycardia, sinus arrest and heart block.
Ranolazine	Ranolazine C _{max} (↑↑), AUC (↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of ranolazine-related adverse reactions, such as QT prolongation and renal failure.
Contraceptives*		
Dienogest Ulipristal	Dienogest C _{max} (↑), AUC (↑↑) ^a Ulipristal C _{max} (↑↑), AUC (↑↑↑) ^a	Use with caution, monitor for contraceptive adverse reactions ^c , refer to the dienogest/ulipristal label for specific actions to be taken.
Diuretics		
Eplerenone	Eplerenone C _{max} (↑), AUC (↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of eplerenone-related adverse reactions, such as hyperkalemia and hypotension.
Finerenone	Finerenone C _{max} (↑↑), AUC (↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of finerenone-related adverse reactions ^c .
Gastrointestinal Drugs		
Aprepitant Loperamide Netupitant	Aprepitant AUC (↑↑↑) ^a Loperamide C _{max} ↑↑, AUC ↑↑ Netupitant C _{max} (↑), AUC (↑↑) ^a	Use with caution, monitor for aprepitant/loperamide/netupitant adverse reactions ^c . Dose reduction of aprepitant/loperamide/ may be necessary. Refer to the netupitant label for specific actions to be taken.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
Cisapride	Cisapride conc increase (extent unknown) ^{a,b}	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of cisapride-related adverse reactions, such as serious cardiovascular events including QT prolongation, serious ventricular arrhythmias and TdP.
Domperidone	Domperidone C _{max} ↑↑, AUC ↑↑	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of domperidone-related adverse reactions, such as serious ventricular arrhythmias and sudden cardiac death.
Drugs that reduce gastric acidity	Itraconazole: C _{max} ↓↓, AUC ↓↓	Use with caution: Drugs that reduce gastric acidity: e.g. acid neutralizing medicines such as aluminum hydroxide, or acid secretion suppressors such as H ₂ -receptor antagonists and proton pump inhibitors. When co-treatment with acid neutralizing medicines (e.g. aluminum hydroxide) these should be administered at least 2 hours before or 2 hours after the intake of SPORAL capsules. (See 4.4 Special warning and precautions.)
Naloxegol	Naloxegol C _{max} (↑↑↑), AUC (↑↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of naloxegol-related adverse reactions, such as opioid withdrawal symptoms.
<i>Saccharomyces boulardii</i>	<i>S. boulardii</i> colonization decrease (extent unknown)	Not recommended during and for 2 weeks after treatment with itraconazole. <i>S. boulardii</i> efficacy may be reduced.
Immunosuppressants		
Budesonide Ciclesonide Cyclosporine Dexamethasone	Budesonide (inhalation) C _{max} ↑, AUC ↑↑; Budesonide (other form.) conc increase (extent unknown) ^{a,b}	Use with caution, monitor for immunosuppressant adverse reactions ^c , Dose reduction of the immunosuppressant drug may be necessary.

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
Fluticasone Methylprednisolone Tacrolimus Temsirolimus	Ciclesonide (inhalation) C_{max} ($\uparrow\uparrow$), AUC ($\uparrow\uparrow$) ^a Cyclosporine (iv) conc increase \leftrightarrow to \uparrow^b Cyclosporine (other form.) conc increase (extent unknown) ^{a,b} Dexamethasone C_{max} \leftrightarrow (iv) \uparrow (oral), AUC $\uparrow\uparrow$ (iv, oral) Fluticasone (inhalation) conc increase $\uparrow\uparrow^b$ Fluticasone (nasal) conc increase (\uparrow) ^{a,b} Methylprednisolone (oral) C_{max} \uparrow to $\uparrow\uparrow$, AUC $\uparrow\uparrow$ Methylprednisolone (iv) AUC $\uparrow\uparrow$ Tacrolimus (iv) conc increase \uparrow^b Tacrolimus (oral) C_{max} ($\uparrow\uparrow$), AUC ($\uparrow\uparrow$) ^a Temsirolimus (iv) C_{max} ($\uparrow\uparrow$), AUC ($\uparrow\uparrow$) ^a	
Everolimus Sirolimus (rapamycin)	Everolimus C_{max} ($\uparrow\uparrow$), AUC ($\uparrow\uparrow\uparrow$) ^a Sirolimus C_{max} ($\uparrow\uparrow$), AUC ($\uparrow\uparrow\uparrow$) ^a	Not recommended during and for 2 weeks after treatment with itraconazole. Increased risk of everolimus/ sirolimus-related adverse reactions ^c .
Voclosporin	Voclosporin C_{max} ($\uparrow\uparrow\uparrow$), AUC ($\uparrow\uparrow\uparrow$) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of voclosporin-related adverse reactions ^c .
Lipid Regulating Drugs		
Atorvastatin	Atorvastatin C_{max} \leftrightarrow to $\uparrow\uparrow$, AUC \uparrow to $\uparrow\uparrow$	Use with caution, monitor for atorvastatin adverse reactions ^c . Dose

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
		reduction of atorvastatin may be necessary.
Lomitapide	Lomitapide C _{max} (↑↑↑↑), AUC (↑↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of lomitapide-related adverse reactions, such as hepatotoxicity and severe gastrointestinal reactions.
Lovastatin Simvastatin	Lovastatin C _{max} ↑↑↑↑, AUC ↑↑↑↑ Simvastatin C _{max} ↑↑↑↑, AUC ↑↑↑↑	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of lovastatin/simvastatin-related adverse reactions, such as myopathy, rhabdomyolysis and liver enzyme abnormalities.
Nonsteroidal Anti-Inflammatory Drugs		
Meloxicam	Meloxicam C _{max} ↓↓, AUC ↓	Use with caution, monitor for reduced efficacy of meloxicam, dose adaption of meloxicam may be necessary.
Respiratory Drugs		
Salmeterol	Salmeterol C _{max} (↑), AUC (↑↑↑↑) ^a	Not recommended during and for 2 weeks after treatment with itraconazole. Increased risk of salmeterol-related adverse reactions ^c .
SSRIs, Tricyclics and Related Antidepressants		
Reboxetine Venlafaxine	Reboxetine C _{max} (↔), AUC (↑) ^a Venlafaxine C _{max} (↑), AUC (↑) ^a	Use with caution, monitor for reboxetine/venlafaxine adverse reactions ^c , dose reduction of reboxetine/venlafaxine may be necessary.
Urologic Drugs		
Avanafil	Avanafil C _{max} (↑↑), AUC (↑↑↑↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk avanafil-related adverse reactions, such as priapism, visual problems and sudden loss of hearing.
Dapoxetine	Dapoxetine C _{max} (↑), AUC (↑) ^a	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk for dapoxetine-related adverse reactions, such as orthostatic hypotension and ocular effects.
Darifenacin Vardenafil	Darifenacin C _{max} (↑↑↑), AUC (↑↑↑ to ↑↑↑↑) ^a Vardenafil C _{max} (↑↑), AUC (↑↑↑↑) ^a	Not recommended during and for 2 weeks after treatment with itraconazole. Increased risk of darifenacin/vardenafil-related adverse reactions ^c .

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
Dutasteride Imidafenacin Oxybutynin Sildenafil (erectile dysfunction) Tadalafil (erectile dysfunction and benign prostatic hyperplasia) Tolterodine Udenafil	Dutasteride conc increase (extent unknown) ^{a,b} Imidafenacin C _{max} ↑, AUC ↑ Oxybutynin conc increase ↑ ^b Sildenafil C _{max} (↑↑), AUC (↑↑ to ↑↑↑↑) ^a Tadalafil C _{max} (↑), AUC (↑↑) ^a Tolterodine C _{max} (↑ to ↑↑), AUC (↑↑) ^a in poor metabolizers of CYP2D6 Udenafil C _{max} (↑), AUC (↑↑) ^a	Use with caution, monitor for urologic drug adverse reactions ^c , dose reduction of the urologic drug may be necessary; refer to the dutasteride label for specific actions to be taken. (For sildenafil and tadalafil, see also <i>Cardiovascular Drugs, Miscellaneous Drugs and other substances.</i>)
Fesoterodine	Fesoterodine C _{max} (↑↑), AUC (↑↑) ^a	Contraindicated in patients with moderate to severe renal or hepatic impairment, during and for 2 weeks after treatment with itraconazole. Increased risk of fesoterodine-related adverse reactions, such as severe anticholinergic effects. Use with caution in other patients: monitor for fesoterodine adverse reactions ^c , dose reduction of fesoterodine may be necessary.
Solifenacin	Solifenacin C _{max} (↑), AUC (↑↑) ^a	Contraindicated in patients with severe renal or moderate to severe hepatic impairment, during and for 2 weeks after treatment with itraconazole. Increased risk of solifenacin-related adverse reactions, such as anticholinergic effects and QT prolongation. Use with caution in other patients, monitor for solifenacin drug adverse reactions ^c , dose reduction of solifenacin may be necessary.
Miscellaneous Drugs and Other Substances		

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
Alitretinoin (oral) Cabergoline Cannabinoids Cinacalcet	Alitretinoin C_{max} (↑), AUC (↑) ^a Cabergoline C_{max} (↑↑), AUC (↑↑) ^a Cannabinoids conc increase, extent unknown but likely (↑↑) ^a Cinacalcet C_{max} (↑↑), AUC (↑↑) ^a	Use with caution, monitor for alitretinoin/cabergoline/cannabinoids/cinacalcet drug adverse reactions ^c , dose reduction of alitretinoin/cabergoline/cannabinoids/cinacalcet may be necessary.
Valbenazine	Valbenazine C_{max} (↑), AUC (↑↑) ^a	Use with caution, monitor for valbenazine-related adverse reactions ^c , dose reduction of valbenazine is necessary.
Colchicine	Colchicine C_{max} (↑), AUC (↑↑) ^a	Contraindicated in patients with renal or hepatic impairment, during and for 2 weeks after treatment with itraconazole. Increased risk of colchicine-related adverse reactions, such as decreased cardiac output, cardiac arrhythmias, respiratory distress and bone marrow depression. Not recommended in other patients, during and for 2 weeks after treatment with itraconazole. Increased risk of colchicine-related adverse reactions ^c .
Eliglustat	CYP2D6 EMs: Eliglustat C_{max} (↑↑), AUC (↑↑) ^a Higher increases are expected in CYP2D6 IMs/PMs and upon coadministration with a CYP2D6 inhibitor.	Contraindicated in CYP2D6 EMs taking a strong or moderate CYP2D6 inhibitor / CYP2D6 IMs and PMs, during and for 2 weeks after treatment with itraconazole. Increased risk of eliglustat-related adverse reactions such as prolongation of the PR, QTc, and/or QRS cardiac interval, and cardiac arrhythmias. Use with caution in CYP2D6 EMs, monitor for eliglustat adverse reactions ^c , dose reduction of eliglustat may be necessary.
Ergot alkaloids	Ergot alkaloids conc increase (extent unknown) ^{a,b}	Contraindicated during and for 2 weeks after treatment with itraconazole. Increased risk of ergot alkaloid-related adverse reactions, such as ergotism. (see also <i>Antimigraine Drugs</i>)
Galantamine	Galantamine C_{max} (↑), AUC (↑) ^a	Use with caution, monitor for galantamine adverse reactions ^c . Dose

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
		reduction of galantamine may be necessary.
Ivacaftor	Ivacaftor C _{max} (↑↑), AUC (↑↑↑) ^a	Use with caution, monitor for ivacaftor adverse reactions ^c , dose reduction of ivacaftor may be necessary.
Lumacaftor/Ivacaftor	Ivacaftor C _{max} (↑↑), AUC (↑↑) ^a Lumacaftor C _{max} (↔), AUC (↔) ^a Itraconazole conc decrease, extent unknown but likely ↓↓↓	Not recommended from 2 weeks before, during and for 2 weeks after treatment with itraconazole. Itraconazole efficacy may be reduced and increased risk of ivacaftor-related adverse reactions ^c .
Vasopressin Receptor Antagonists		
Conivaptan Tolvaptan	Conivaptan C _{max} (↑↑), AUC (↑↑↑↑) ^a Tolvaptan C _{max} (↑↑), AUC (↑↑↑) ^a	Not recommended during and for 2 weeks after treatment with itraconazole. Increased risk of conivaptan/ tolvaptan-related adverse reactions ^c .
Mozavaptan	Mozavaptan C _{max} ↑, AUC ↑↑	Use with caution, monitor for mozavaptan adverse reactions ^c , dose reduction of mozavaptan may be necessary.

*CYP3A4 inhibitors (including itraconazole) may increase systemic contraceptive hormone concentrations.

EMs: extensive metabolizers; IMs: intermediate metabolizers, PMs: poor metabolizers; TdP: Torsade de Pointes

Note:

Average increase:

↑: <100% (i.e. <2-fold);

↑↑: 100-400% (i.e. ≥2-fold to <5-fold);

↑↑↑: 400-900% (i.e. ≥5-fold and <10-fold);

↑↑↑↑: ≥10-fold;

Average decrease:

↓: <40%;

↓↓: 40-80%;

↓↓↓: >80%;

Examples of medicinal products within class	Expected/Potential effect on drug levels (see footnotes for additional info)	Clinical comment (see above for additional info)
No effect: ↔;		

For the effect (middle column) the name of the parent drug is stated, even when the effect is related to the active moiety or the active metabolite of a prodrug.

- ^a For drugs with arrows between brackets, the assessment was based on the mechanism of interaction and clinical drug interaction information with ketoconazole or other strong CYP3A4 inhibitors and/or inhibitors of P-glycoprotein or BCRP, modelling techniques, case reports and/or *in vitro* data. For the other drugs listed, the assessment was based on clinical drug interaction information with itraconazole.
- ^b Pharmacokinetic parameters were not available.
- ^c Please consult the corresponding label for information on drug-related adverse reactions

Pediatric population

Interaction studies have only been performed in adults.

4.6 Fertility, pregnancy and lactation

Pregnancy

SPORAL must not be used during pregnancy except for life-threatening cases where the potential benefit to the mother outweighs the potential harm to the fetus (See section 4.3 *Contraindications*).

In animal studies itraconazole has shown reproduction toxicity (See section 5.3 *Preclinical safety data*).

There is limited information on the use of SPORAL during pregnancy. During post-marketing experience, cases of congenital abnormalities have been reported. These cases included skeletal, genitourinary tract, cardiovascular and ophthalmic malformations as well as chromosomal and multiple malformations. A causal relationship with SPORAL has not been established.

Epidemiological data on exposure to SPORAL during the first trimester of pregnancy – mostly in patients receiving short-term treatment for vulvovaginal candidosis – did not show an increased risk for malformations as compared to control subjects not exposed to any known teratogens. Itraconazole has been shown to cross the placenta in a rat model.

Women of childbearing potential

Women of childbearing potential taking SPORAL capsules should use contraceptive precautions. Highly effective contraception should be continued until the menstrual period following the end of SPORAL therapy.

Breast-feeding

A very small amount of itraconazole is excreted in human milk. The expected benefits of SPORAL capsules therapy should therefore be weighed against the potential risk of breast-feeding. In case of doubt, the patient should not breast-feed.

Fertility

Refer to *Non-Clinical Information for in animal* fertility information relevant to itraconazole.

4.7 Effect on ability to drive and use machine

No studies on the effects on the ability to drive and use machines have been performed. When driving vehicles and operating machinery the possibility of adverse reactions such as dizziness, visual disturbances and hearing loss (See section 4.8 *Undesirable effects.*), which may occur in some instances, must be taken into account.

4.8 Undesirable effects

Throughout this section, adverse reactions are presented. Adverse reactions are adverse events that were considered to be reasonably associated with the use of itraconazole based on the comprehensive assessment of the available adverse event information. A causal relationship with itraconazole cannot be reliably established in individual cases. Further, because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Clinical trial data

The safety of SPORAL capsules was evaluated in 8499 patients who participated in 107 open-label and double-blind clinical trials. Of the 8499 patients treated with SPORAL capsules, 2104 patients were treated with SPORAL capsules during double-blind trials. All 8499 patients received at least one dose of SPORAL capsules for the treatment of dermatomycoses or onychomycosis and provided safety data. Adverse reactions reported for $\geq 1\%$ of patients treated with SPORAL capsules in these clinical trials are shown in Table 1.

Table 1: Adverse Reactions Reported by $\geq 1\%$ of Patients Treated with SPORAL Capsules in 107 Clinical Trials

System Organ Class Adverse Reaction	SPORAL Capsules % (N=8499)
Nervous System Disorders	
Headache	1.6
Gastrointestinal Disorders	
Nausea	1.6
Abdominal pain	1.3

Adverse reactions that occurred in <1% of patients treated with SPORAL capsules in these clinical trials are listed in Table 2.

Table 2: Adverse Reactions Reported by <1% of Patients Treated with SPORAL Capsules in 107 Clinical Trials

System Organ Class
Adverse Reaction
Infections and Infestations
Rhinitis
Sinusitis
Upper respiratory tract infection
Blood and Lymphatic System Disorders
Leukopenia
Immune System Disorders
Hypersensitivity
Nervous System Disorders
Dysgeusia
Hypoesthesia
Paresthesia
Ear and Labyrinth Disorders
Tinnitus
Gastrointestinal Disorders
Constipation
Diarrhea
Dyspepsia
Flatulence
Vomiting
Hepatobiliary Disorders
Hepatic function abnormal
Hyperbilirubinemia
Skin and Subcutaneous Tissue Disorders
Pruritus
Rash
Urticaria
Renal and Urinary Disorders
Pollakiuria
Reproductive System and Breast Disorders
Erectile dysfunction
Menstrual disorder
General Disorders and Administration Site Conditions
Edema

The following is a list of additional adverse reactions associated with itraconazole that have been reported in clinical trials of SPORAL oral solution and/or SPORAL IV, excluding the adverse reaction term "Injection site inflammation" which is specific to the injection route of administration.

Blood and Lymphatic System Disorders: Granulocytopenia, Thrombocytopenia

Immune System Disorders: Anaphylactoid reaction

Metabolism and Nutrition Disorders: Hyperglycemia, Hyperkalemia, Hypokalemia, Hypomagnesemia

Psychiatric Disorders: Confusional state

Nervous System Disorders: Neuropathy peripheral, Dizziness, Somnolence

Cardiac Disorders: Cardiac failure, Left ventricular failure, Tachycardia

Vascular Disorders: Hypertension, Hypotension

Respiratory, Thoracic and Mediastinal Disorders: Pulmonary edema, Dysphonia, Cough

Gastrointestinal Disorders: Gastrointestinal disorder

Hepatobiliary Disorders: Hepatic failure, Hepatitis, Jaundice

Skin and Subcutaneous Tissue Disorders: Rash erythematous, Hyperhidrosis

Musculoskeletal and Connective Tissue Disorders: Myalgia, Arthralgia

Renal and Urinary Disorders: Renal impairment, Urinary incontinence

General Disorders and Administration Site Conditions: Generalized edema, Face edema, Chest pain, Pyrexia, Pain, Fatigue, Chills

Investigations: Alanine aminotransferase increased, Aspartate aminotransferase increased, Blood alkaline phosphatase increased, Blood lactate dehydrogenase increased, Blood urea increased, Gamma-glutamyltransferase increased, Hepatic enzyme increased, Urine analysis abnormal

Pediatrics

The safety of SPORAL capsules was evaluated in 165 pediatric patients aged 1 to 17 years who participated in 14 clinical trials (4 double-blind, placebo controlled trials; 9 open-label trials; and 1 trial had an open-label phase followed by a double-blind phase). These patients received at least one dose of SPORAL capsules for the treatment of fungal infections and provided safety data.

Based on pooled safety data from these clinical trials, the commonly reported adverse reactions in pediatric patients were Headache (3.0%), Vomiting (3.0%), Abdominal pain (2.4%), Diarrhea (2.4%), Hepatic function abnormal (1.2%), Hypotension (1.2%), Nausea (1.2%), and Urticaria (1.2%). In general, the nature of adverse reactions in pediatric patients is similar to that observed in adult subjects, but the incidence is higher in the pediatric patients.

Postmarketing data

In addition to the adverse reactions reported during clinical studies and listed above, the following adverse reactions have been reported during postmarketing experience

(Table 3). In each table, the frequencies are provided according to the following convention:

Very common	≥ 1/10
Common	≥ 1/100 and < 1/10
Uncommon	≥ 1/1000 and < 1/100
Rare	≥ 1/10000 and < 1/1000
Very rare	< 1/10000, including isolated reports.

In Table 3, adverse reactions are presented by frequency category based on spontaneous reporting rates.

Table 3: Adverse Reactions Identified During Post-Marketing Experience with SPORAL by Frequency Category Estimated from Spontaneous Reporting Rates

Immune System Disorders	
<i>Very rare</i>	Serum sickness, Angioneurotic edema, Anaphylactic reaction
Endocrine Disorders	
<i>Very rare</i>	Pseudoaldosteronism
Metabolism and Nutrition Disorders	
<i>Very rare</i>	Hypertriglyceridemia
Nervous System Disorders	
<i>Very rare</i>	Tremor
Eye Disorders	
<i>Very rare</i>	Visual disturbances (including diplopia and vision blurred)
Ear and Labyrinth Disorders	
<i>Very rare</i>	Transient or permanent hearing loss
Cardiac Disorders	
<i>Very rare</i>	Congestive heart failure, Bradycardia
Respiratory, Thoracic and Mediastinal Disorders	
<i>Very rare</i>	Dyspnea
Gastrointestinal Disorders	
<i>Very rare</i>	Pancreatitis
Hepatobiliary Disorders	
<i>Very rare</i>	Serious hepatotoxicity (including some cases of fatal acute liver failure)
Skin and Subcutaneous Tissue Disorders	
<i>Very rare</i>	Toxic epidermal necrolysis, Stevens-Johnson syndrome, Acute generalized exanthematous pustulosis, Erythema multiforme, Exfoliative dermatitis, Leukocytoclastic vasculitis, Alopecia, Photosensitivity
Investigations	
<i>Very rare</i>	Blood creatine phosphokinase increased

4.9 Overdose

Symptoms and signs

In general, adverse events reported with overdose have been consistent with those reported for itraconazole use (See section 4.8 *Undesirable effects*).

Treatment

In the event of an overdose, supportive measures should be employed.

It is advisable to contact a poison control center to determine the latest recommendations for the management of an overdose.

Itraconazole cannot be removed by hemodialysis.

No specific antidote is available.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic Properties

Pharmacotherapeutic group: Antimycotic for systemic use, triazole and tetrazole derivatives, ATC code: J02A C02.

Mechanism of action

In vitro studies have demonstrated that itraconazole impairs the synthesis of ergosterol in fungal cells. Ergosterol is a vital cell membrane component in fungi. Impairment of its synthesis ultimately results in an antifungal effect.

Pharmacokinetic (PK)/Pharmacodynamic (PD) relationship

The PK/PD relationship for itraconazole, and for triazoles in general, is poorly understood.

Pharmacodynamic effects

Microbiology

Itraconazole, a triazole derivative, has a broad spectrum of activity.

For itraconazole, interpretive breakpoints have not been established by CLSI for *Candida* spp. and the filamentous fungi.

EUCAST breakpoints for itraconazole have been established for *Aspergillus flavus*, *A. fumigatus*, *A. nidulans* and *A. terreus*, and are as follows: susceptible ≤ 1 mg/L, resistant > 1 mg/L. EUCAST breakpoints for itraconazole have been established for *Candida albicans* and *C. dubliniensis*, and are as follows: susceptible ≤ 0.06 mg/L, resistant > 0.06 mg/L. EUCAST breakpoints for itraconazole have been established for *Candida parapsilosis* and *C. tropicalis*, and are as follows: susceptible ≤ 0.125 mg/L, resistant > 0.125 mg/L. Interpretive breakpoints have not been established by EUCAST for *Candida glabrata*, *C. krusei*, *C. guilliermondii*, *Cryptococcus neoformans*, *Aspergillus niger*, and Non-species related breakpoints for *Candida* and *Aspergillus*.

In vitro studies demonstrate that itraconazole inhibits the growth of a broad range of fungi pathogenic for humans at concentrations usually $\leq 1 \mu\text{g/mL}$. These include:

Aspergillus spp., *Blastomyces dermatitidis*, *Cladosporium* spp., *Coccidioides immitis*, *Cryptococcus neoformans*, *Geotrichum* spp., *Histoplasma* spp., including *H. capsulatum*, *Paracoccidioides brasiliensis*, *Talaromyces* (formerly *Penicillium*) *marneffeii*, *Sporothrix schenckii* and *Trichosporon* spp. Itraconazole also displayed activity *in vitro* against *Epidermophyton floccosum*, *Fonsecaea* spp., *Malassezia* spp., *Microsporium* spp., *Pseudallescheria boydii*, *Trichophyton* spp. and various other yeasts and fungi.

The principal fungus types that are not inhibited by itraconazole are Zygomycetes (e.g. *Rhizopus* spp., *Rhizomucor* spp., *Mucor* spp. and *Absidia* spp.), *Fusarium* spp., *Scedosporium* spp. and *Scopulariopsis* spp.

Azole resistance appears to develop slowly and is often the result of several genetic mutations. Mechanisms that have been described are overexpression of ERG11, which encodes the target enzyme 14 α -demethylase, point mutations in ERG11 that lead to decreased target affinity and/or transporter overexpression resulting in increased efflux. Cross-resistance between members of the azole class has been observed within *Candida* spp., although resistance to one member of the class does not necessarily confer resistance to other azoles. Itraconazole-resistant strains of *Aspergillus fumigatus* have been reported.

5.2 Pharmacokinetic Properties

General pharmacokinetic characteristics

Peak plasma concentrations of itraconazole are reached within 2 to 5 hours following oral administration. As a consequence of non-linear pharmacokinetics, itraconazole accumulates in plasma during multiple dosing. Steady-state concentrations are generally reached within about 15 days, with C_{max} values of 0.5 $\mu\text{g/mL}$, 1.1 $\mu\text{g/mL}$ and 2.0 $\mu\text{g/mL}$ after oral administration of 100 mg once daily, 200 mg once daily and 200 mg b.i.d., respectively. The terminal half-life of itraconazole generally ranges from 16 to 28 hours after single dose and increases to 34 to 42 hours with repeated dosing. Once treatment is stopped, itraconazole plasma concentrations decrease to an almost undetectable concentration within 7 to 14 days, depending on the dose and duration of treatment. Itraconazole mean total plasma clearance following intravenous administration is 278 mL/min. Itraconazole clearance decreases at higher doses due to saturable hepatic metabolism.

Absorption

Itraconazole is rapidly absorbed after oral administration. Peak plasma concentrations of the unchanged drug are reached within 2 to 5 hours following an oral capsule dose. The observed absolute oral bioavailability of itraconazole is about 55%. Oral bioavailability is maximal when the capsules are taken immediately after a full meal.

Absorption of itraconazole capsules is reduced in subjects with reduced gastric acidity, such as subjects taking medications known as gastric acid secretion suppressors (e.g., H₂-receptor antagonists, proton pump inhibitors) or subjects with achlorhydria caused by certain diseases (see section 4.4 *Special warning and precautions*, and section 4.5

Interaction with other medicinal products and other forms of interactions). Absorption of itraconazole under fasted conditions in these subjects is increased when SPORAL capsules are administered with an acidic beverage (such as a non-diet cola). When SPORAL capsules were administered as a single 200-mg dose under fasted conditions with non-diet cola after ranitidine pretreatment, a H₂-receptor antagonist, itraconazole absorption was comparable to that observed when SPORAL capsules were administered alone (See section 4.5 *Interaction with other medicinal products and other forms of interactions*).

Itraconazole exposure is lower with the capsule formulation than with the oral solution when the same dose of drug is given (see section 4.4 *Special warning and precautions*).

Distribution

Most of the itraconazole in plasma is bound to protein (99.8%), with albumin being the main binding component (99.6% for the hydroxy-metabolite). It has also a marked affinity for lipids. Only 0.2% of the itraconazole in plasma is present as free drug. Itraconazole is distributed in a large apparent volume in the body (> 700 L), suggesting extensive distribution into tissues. Concentrations in lung, kidney, liver, bone, stomach, spleen and muscle were found to be two to three times higher than corresponding concentrations in plasma, and the uptake into keratinous tissues, skin in particular, up to four times higher. Concentrations in the cerebrospinal fluid are much lower than in plasma, but efficacy has been demonstrated against infections present in the cerebrospinal fluid.

Metabolism

Itraconazole is extensively metabolized by the liver into a large number of metabolites. *In vitro* studies have shown that CYP3A4 is the major enzyme involved in the metabolism of itraconazole. The main metabolite is hydroxy-itraconazole, which has *in vitro* antifungal activity comparable to itraconazole; trough plasma concentrations of this metabolite are about twice those of itraconazole.

Excretion

Itraconazole is excreted mainly as inactive metabolites in urine (35%) and in feces (54%) within one week of an oral solution dose. Renal excretion of itraconazole and the active metabolite hydroxy-itraconazole account for less than 1% of an intravenous dose. Based on an oral radiolabeled dose, fecal excretion of unchanged drug ranges from 3% to 18% of the dose.

As re-distribution of itraconazole from keratinous tissues appears to be negligible, elimination of itraconazole from these tissues is related to epidermal regeneration. Contrary to plasma, the concentration in skin persists for 2 to 4 weeks after discontinuation of a 4-week treatment and in nail keratin – where itraconazole can be detected as early as 1 week after start of treatment – for at least six months after the end of a 3-month treatment period.

Special populations

Hepatic impairment

Itraconazole is predominantly metabolized in the liver. A pharmacokinetic study was conducted in 6 healthy and 12 cirrhotic subjects who were administered a single 100-mg dose of itraconazole as a capsule. A statistically significant reduction in mean C_{max} (47%) and a twofold increase in the elimination half-life (37 ± 17 hours vs. 16 ± 5 hours) of itraconazole were noted in cirrhotic subjects compared with healthy subjects. However, overall exposure to itraconazole, based on AUC, was similar in cirrhotic patients and in healthy subjects. Data are not available in cirrhotic patients during long-term use of itraconazole (See section 4.2 *Posology and method of administration*, and section 4.4 *Special warning and precautions for use*).

Renal impairment

Limited data are available on the use of oral itraconazole in patients with renal impairment. A pharmacokinetic study using a single 200-mg dose of itraconazole (four 50-mg capsules) was conducted in three groups of patients with renal impairment (uremia: n=7; hemodialysis: n=7; and continuous ambulatory peritoneal dialysis: n=5). In uremic subjects with a mean creatinine clearance of $13 \text{ mL/min} \times 1.73 \text{ m}^2$, the exposure, based on AUC, was slightly reduced compared with normal population parameters. This study did not demonstrate any significant effect of hemodialysis or continuous ambulatory peritoneal dialysis on the pharmacokinetics of itraconazole (T_{max} , C_{max} , and AUC_{0-8h}). Plasma concentration-versus-time profiles showed wide intersubject variation in all three groups.

After a single intravenous dose, the mean terminal half-lives of itraconazole in patients with mild (defined in this study as CrCl 50-79 mL/min), moderate (defined in this study as CrCl 20-49 mL/min), and severe renal impairment (defined in this study as CrCl <20 mL/min) were similar to that in healthy subjects, (range of means 42-49 hours vs 48 hours in renally impaired patients and healthy subjects, respectively.) Overall exposure to itraconazole, based on AUC, was decreased in patients with moderate and severe renal impairment by approximately 30% and 40%, respectively, as compared with subjects with normal renal function.

Data are not available in renally impaired patients during long-term use of itraconazole. Dialysis has no effect on the half-life or clearance of itraconazole or hydroxy-itraconazole (See also section 4.2 *Posology and method of administration*, and section 4.4 *Special warning and precautions for use*).

Pediatrics

Limited pharmacokinetic data are available on the use of itraconazole in the pediatric population. Clinical pharmacokinetic studies in children and adolescents aged between 5 months and 17 years were performed with itraconazole capsules, oral solution or intravenous formulation. Individual doses with the capsule and oral solution formulation ranged from 1.5 to 12.5 mg/kg/day, given as once-daily or twice-daily administration. The intravenous formulation was given either as a 2.5 mg/kg single infusion, or a 2.5 mg/kg infusion given once daily or twice daily. For the same daily dose, twice daily dosing compared to single daily dosing yielded peak and trough concentrations comparable to adult single daily dosing. No significant age dependence was observed for itraconazole AUC and total body clearance, while weak associations between age and itraconazole distribution volume, C_{max} and terminal elimination rate were noted. Itraconazole apparent clearance and distribution volume seemed to be related to weight.

5.3 Preclinical safety data

Itraconazole has been tested in a standard battery of non-clinical safety studies.

Acute oral toxicity studies with itraconazole in mice, rats, guinea pigs and dogs indicate a wide safety margin (4- to 16-fold of Maximum Recommended Human Dose [MRHD] of 400 mg/day based on mg/m²/day). Sub (chronic) oral toxicity studies in rats and dogs revealed several target organs or tissues: adrenal cortex, liver and mononuclear phagocyte system as well as disorders of the lipid metabolism presenting as xanthoma cells in various organs.

At high doses of 40 and 80 mg/kg/day in rats (1- and 2-fold of MRHD based on mg/m²/day, histological investigations of adrenal cortex showed a reversible swelling with cellular hypertrophy of the zona reticularis and fasciculata, which was sometimes associated with a thinning of the zona glomerulosa. Reversible hepatic changes were found at 40 and 160 mg/kg/day (1- and 4-fold of MRHD based on mg/m²/day). Slight changes were observed in the sinusoidal cells and vacuolation of the hepatocytes, the latter indicating cellular dysfunction, but without visible hepatitis or hepatocellular necrosis. Histological changes of the mononuclear phagocyte system were mainly characterized by macrophages with increased proteinaceous material in various parenchymal tissues.

A global lower bone mineral density was observed in juvenile dogs after chronic itraconazole administration. No toxicity was observed up to 20 mg/kg/day (2-fold of MRHD based on mg/m²/day).

In three toxicology studies using rats, itraconazole induced bone defects. The induced defects included reduced bone plate activity, thinning of the zona compacta of the large bones, and an increased bone fragility.

Carcinogenicity and Mutagenicity

Itraconazole is not a primary carcinogen in rats up to 13 mg/kg/day (males) and 52 mg/kg/day (females), or in mice up to 80 mg/kg/day (1-fold of MRHD based on mg/m²/day). In male rats at 25 mg/kg/day dose (0.6-fold of MRHD based on mg/m²/day), however, there was a higher incidence of soft-tissue sarcoma, which is attributed to the increase in non-neoplastic, chronic inflammatory reactions of the connective tissue as a consequence of raised cholesterol levels and cholesterosis in connective tissue.

There are no indications of a mutagenic potential of itraconazole.

Reproductive Toxicology

Itraconazole was found to cause a dose-related increase in maternal toxicity, embryotoxicity, and teratogenicity in rats at 40 and 160 mg/kg/day (1- and 4-fold of MRHD based on mg/m²/day) and mice at 80 and 160 mg/kg/day (1- and 2-fold of MRHD based on mg/m²/day). In rats, the teratogenicity consisted of major skeletal defects; in mice, it consisted of encephaloceles and macroglossia. The observed skeletal malformation in rats may be due to maternal toxicity. No teratogenic effects were found in rabbits up to 80 mg/kg/day dose (4-fold of MRHD based on mg/m²/day).

Fertility

There is no evidence of a primary influence on fertility under treatment with itraconazole.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

The inactive ingredients of the capsules are hypromellose, macrogol, and sugar spheres.

6.2 Incompatibilities

None known.

6.3 Shelf life

See expiry date on the outer pack.

6.4 Special precautions for storage

Store below 30° C.

Keep out of the sight and reach of children.

6.5 Nature and contents of container

SPORAL is available as blue opaque cap and pink transparent body, containing 100 mg of itraconazole in a pellet formulation, supplied in blister packs with 1x4's.

Instructions for Use and Handling

Not applicable.

7. MARKETING AUTHORISATION HOLDER

Imported by

Janssen-Cilag Ltd., Bangkok, Thailand

To report Suspected Adverse Reactions, please contact us at aepqcjacth@its.jnj.com

For any product information, please contact us at medinfosea@its.jnj.com

Manufactured by

Janssen Cilag SpA., Latina, Italy

8. MARKETING AUTHORISATION NUMBER

1C 15010/64

9. DATE OF FIRST AUTHORISATION

18 March 2021

10. DATE OF REVISION OF THE TEXT

CCDS version 27-Mar-2024