

GLUCOSE 50%

กลูโคส 50%

Composition : Each 100 mL contains:-

Dextrose anhydrous 50% w/v

Indication :

1. Used in patient with low blood sugar (Hypoglycemic)
 - Hypoglycemic emergencies
 - Insulin or suspected insulin coma, hypoglycemic shock in patient who takes an overdose of hypoglycemic agents.
2. Used in patients with hyperkalemia.

Dosage and administration :

- Adults and children with hypoglycemic by insulin; intravenous injection 20–50 mL of as directed by the physician.
- Hypoglycemic emergencies; intravenous injection 50 mL or as directed by the physician.
- Hyperkalemia

Children : 0.5–1 g/kg combined with regular insulin 1 unit for every 4–5 g dextrose given; infuse over 2 hours.

Adults : 25–50 g dextrose combined with 10 units regular insulin administered over 30–60 minutes.

Precaution :

1. Intravenous only because subcutaneous may cause tissue necrosis.
2. Should intravenous injection slowly because may cause water lose of tissue.
3. Should not be administered with blood by infusion set.

Storage condition :

Store below 30°C

Manufactured by :

ABLE MEDICAL COMPANY LIMITED

111 Moo 9, Nong Son, Chiang Yuen, Mahasarakham 44160, Thailand