



## CIBINQO™

### 1. NAME OF THE MEDICINAL PRODUCT

#### 1.1 Product Name

CIBINQO 50 mg

CIBINQO 100 mg

CIBINQO 200 mg

#### 1.2 Strength

50 mg, 100 mg and 200 mg

#### 1.3 Pharmaceutical Dosage Form

Film-coated tablets.

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

#### 2.1 Qualitative Declaration

Active ingredient: abrocitinib.

#### 2.2 Quantitative Declaration

Each CIBINQO film-coated tablet contains 50 mg, 100 mg or 200 mg of abrocitinib.

### *Excipient with known effect*

Each 50 mg, 100 mg or 200 mg film-coated tablet contains 1.37 mg, 2.73 mg or 5.46 mg of lactose monohydrate.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Film-coated tablet.

**CIBINQO 50 mg film-coated tablets:** Oval pink film-coated tablet debossed with “PFE” on one side and “ABR 50” on the other.

**CIBINQO 100 mg film-coated tablets:** Round pink film-coated tablet debossed with “PFE” on one side and “ABR 100” on the other.

**CIBINQO 200 mg film-coated tablets:** Oval pink film-coated tablet debossed with “PFE” on one side and “ABR 200” on the other.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic Indications**

CIBINQO is indicated for the treatment of moderate-to-severe atopic dermatitis in adults and adolescents 12 years and older who are candidates for systemic therapy.

### **4.2. Posology and Method of Administration**

Treatment should be initiated and supervised by a healthcare professional experienced in the diagnosis and treatment of atopic dermatitis.

#### **Posology**

The recommended starting dose is 100 mg or 200 mg once daily based on individual patient characteristics:

- A starting dose of 100 mg once daily is recommended for patients at higher risk of venous thromboembolism (VTE), major adverse cardiovascular event (MACE) and malignancy (see section 4.4). If the patient does not respond adequately to 100 mg once daily, the dose can be increased to 200 mg once daily.
- A dose of 200 mg once daily may be appropriate for patients who are not at higher risk of VTE, MACE and malignancy with high disease burden or for patients with an inadequate response to 100 mg once daily. Upon disease control, dose should be decreased to 100 mg once daily. If disease control is not maintained after dose reduction, re-treatment with 200 mg once daily can be considered. In adolescents (12 years to 17 years of age), weighing 25 kg to <59 kg, a starting dose of 100 mg once a day is recommended. If the patient does not respond adequately to 100 mg once daily, the dose can be increased to 200 mg once daily. In adolescents weighing at least 59 kg, a starting dose of 100 mg or 200 mg once daily may be appropriate.

The lowest effective dose for maintenance should be considered.

Discontinuation of treatment should be considered in patients who show no evidence of therapeutic benefit after 24 weeks.

CIBINQO can be used with or without medicated topical therapies for atopic dermatitis.

### **Laboratory monitoring**

**Table 1. Laboratory measures and monitoring guidance**

<b>Laboratory measures</b>	<b>Monitoring guidance</b>	<b>Action</b>
Complete blood count including Platelet Count, Absolute Lymphocyte Count (ALC), Absolute Neutrophil Count (ANC) and Haemoglobin (Hb)	Before treatment initiation, 4 weeks after initiation and thereafter according to routine patient management.	Platelets: Treatment should be discontinued if platelet counts are $<50 \times 10^3/\text{mm}^3$ .
		ALC: Treatment should be interrupted if ALC is $<0.5 \times 10^3/\text{mm}^3$ and may be restarted once ALC returns above this value. Treatment should be discontinued if confirmed.
		ANC: Treatment should be interrupted if ANC is $<1 \times 10^3/\text{mm}^3$ and may be restarted once ANC returns above this value.
		Hb: Treatment should be interrupted if Hb is $<8 \text{ g/dL}$ and may be restarted once Hb returns above this value.
Lipid parameters	Before treatment initiation, 4 weeks after initiation and thereafter according to the patient's risk for cardiovascular disease and clinical guidelines for hyperlipidaemia.	Patients should be monitored according to clinical guidelines for hyperlipidaemia.

### **Treatment initiation**

Treatment should not be initiated in patients with a platelet count  $<150 \times 10^3/\text{mm}^3$ , an absolute lymphocyte count (ALC)  $<0.5 \times 10^3/\text{mm}^3$ , an absolute neutrophil count (ANC)  $<1.2 \times 10^3/\text{mm}^3$  or who have a haemoglobin value  $<10 \text{ g/dL}$  (see section 4.4).

### ***Dose interruption***

If a patient develops a serious infection, sepsis or opportunistic infection, dose interruption should be considered until the infection is controlled (see section 4.4).

Interruption of dosing may be needed for management of laboratory abnormalities as described in Table 1.

### ***Missed doses***

If a dose is missed, patients should be advised to take the dose as soon as possible unless it is less than 12 hours before the next dose, in which case the patient should not take the missed dose. Thereafter, dosing should be resumed at the regular scheduled time.

### ***Interactions***

In patients receiving dual strong inhibitors of CYP2C19 and moderate inhibitors of CYP2C9, or strong inhibitors of CYP2C19 alone (e.g., fluvoxamine, fluconazole, fluoxetine and ticlopidine), the recommended dose should be reduced by half to 100 mg or 50 mg once daily (see section 4.5).

Treatment is not recommended concomitantly with moderate or strong inducers of CYP2C19/CYP2C9 enzymes (e.g., rifampicin, apalutamide, efavirenz, enzalutamide, phenytoin) (see section 4.5).

In patients receiving acid reducing agents (e.g. antacids, proton pump inhibitors and H2 receptor antagonists), 200 mg once daily dose of abrocitinib should be considered (see section 4.5).

### ***Special populations***

#### ***Renal impairment***

No dose adjustment is required in patients with mild renal impairment, i.e., estimated glomerular filtration rate (eGFR) of 60 to <90 mL/min.

In patients with moderate (eGFR 30 to <60 mL/min) renal impairment, the recommended dose of

abrocitinib should be reduced by half to 100 mg or 50 mg once daily (see section 5.2).

In patients with severe (eGFR <30 mL/min) renal impairment, 50 mg once daily is the recommended starting dose. The maximum daily dose is 100 mg (see section 5.2).

Abrocitinib has not been studied in patients with end-stage renal disease (ESRD) on renal replacement therapy.

### ***Hepatic impairment***

No dose adjustment is required in patients with mild (Child Pugh A) or moderate (Child Pugh B) hepatic impairment. Abrocitinib is contraindicated to patients with severe (Child Pugh C) hepatic impairment (see section 4.3).

### ***Elderly***

For patients 65 years of age and older, the recommended dose is 100 mg once daily (see section 4.4).

### ***Paediatric population***

The safety and efficacy of CIBINQO in children under 12 years of age have not yet been established. No data are available.

### **Method of administration**

This medicinal product is to be taken orally once daily with or without food at approximately the same time each day.

In patients who experience nausea, taking tablets with food may improve nausea.

Tablets should be swallowed whole with water and should not be split, crushed or chewed because these methods have not been studied in clinical trials.

### 4.3. Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Active serious systemic infections, including tuberculosis (TB) (see section 4.4).
- Severe hepatic impairment (see section 4.2).
- Pregnancy and breast-feeding (see section 4.6).

### 4.4. Special Warnings and Precautions for Use

Abrocitinib should only be used if no suitable treatment alternatives are available in patients:

- 65 years of age and older;
- patients with history of atherosclerotic cardiovascular disease or other cardiovascular risk factors (such as current or past long-time smokers);
- patients with malignancy risk factors (e.g. current malignancy or history of malignancy)

#### Infections/serious infections

Serious infections have been reported in patients receiving abrocitinib. The most frequent serious infections in clinical studies were herpes simplex, herpes zoster, and pneumonia (see section 4.8).

As there is a higher incidence of infections in the elderly and in the diabetic populations in general, caution should be used when treating the elderly and patients with diabetes. In patients 65 years of age and older abrocitinib should only be used if no suitable treatment alternatives are available (see section 4.2).

Treatment must not be initiated in patients with an active, serious systemic infection (see section 4.3).

Risks and benefits of treatment prior to initiating abrocitinib should be considered for patients:

- with chronic or recurrent infection
- who have been exposed to TB
- with a history of a serious or an opportunistic infection
- who have resided or travelled in areas of endemic TB or endemic mycoses; or
- with underlying conditions that may predispose them to infection.

Patients should be closely monitored for the development of signs and symptoms of infection during and after treatment with abrocitinib. A patient who develops a new infection during treatment should undergo prompt and complete diagnostic testing and appropriate antimicrobial therapy should be initiated. The patient should be closely monitored and therapy should be temporarily interrupted if the patient is not responding to standard therapy.

### ***Tuberculosis***

Tuberculosis was observed in clinical studies with abrocitinib. Patients should be screened for TB before starting treatment and yearly screening for patients in highly endemic areas for TB should be considered. Abrocitinib must not be given to patients with active TB (see section 4.3). For patients with a new diagnosis of latent TB or prior untreated latent TB, preventive therapy for latent TB should be started prior to initiation of treatment.

### ***Viral reactivation***

Viral reactivation, including herpes virus reactivation (e.g. herpes zoster, herpes simplex), was reported in clinical studies (see section 4.8). The rate of herpes zoster infections was higher in patients who were treated with 200 mg, 65 years of age and older, with a medical history of herpes zoster, with a confirmed ALC  $<1 \times 10^3/\text{mm}^3$  prior to the event and patients with severe atopic dermatitis at baseline (see section 4.8). If a patient develops herpes zoster, temporary interruption of treatment should be considered until the episode resolves.

Screening for viral hepatitis should be performed in accordance with clinical guidelines before starting therapy and during therapy. Patients with evidence of active hepatitis B or hepatitis C (positive hepatitis C PCR) infection were excluded from clinical studies (see section 5.2). Patients who were hepatitis B surface antigen negative, hepatitis B core antibody positive, and hepatitis B surface antibody positive had testing for hepatitis B virus (HBV) DNA. Patients who had HBV DNA above the lower limit of quantification (LLQ) were excluded. Patients who had HBV DNA negative or below LLQ could initiate treatment; such patients had HBV DNA monitored. If HBV DNA is detected, a liver specialist should be consulted.

### **Vaccination**



No data are available on the response to vaccination in patients receiving abrocitinib. Use of live, attenuated vaccines should be avoided during or immediately prior to treatment. Prior to initiating treatment with this medicinal product, it is recommended that patients be brought up to date with all immunisations, including prophylactic herpes zoster vaccinations, in agreement with current immunisation guidelines.

### **Venous thromboembolism (VTE)**

Events of deep venous thrombosis (DVT) and pulmonary embolism (PE) have been reported in patients receiving abrocitinib (see section 4.8).

In a large randomized active-controlled study of tofacitinib (another JAK inhibitor) in rheumatoid arthritis patients 50 years and older with at least one additional cardiovascular risk factor, a dose dependent higher rate of VTE including deep venous thrombosis (DVT) and pulmonary embolism (PE) was observed with tofacitinib compared to TNF inhibitors.

A higher rate of VTE was observed with abrocitinib 200 mg compared to abrocitinib 100 mg.

In patients with cardiovascular or malignancy risk factors (see also section 4.4 “Major adverse cardiovascular events (MACE)” and “Malignancy”) abrocitinib should only be used if no suitable treatment alternatives are available.

In patients with known VTE risk factors other than cardiovascular or malignancy risk factors, abrocitinib should be used with caution. VTE risk factors other than cardiovascular or malignancy risk factors include previous VTE, patients undergoing major surgery, immobilisation, use of combined hormonal contraceptives or hormone replacement therapy, inherited coagulation disorder.

Patients should be re-evaluated periodically during abrocitinib treatment to assess for changes in VTE risk.

Promptly evaluate patients with signs and symptoms of VTE and discontinue abrocitinib in patients with suspected VTE, regardless of dose.

### **Major adverse cardiovascular events (MACE)**

Events of MACE have been observed in patients taking abrocitinib.

In a large randomized active-controlled study of tofacitinib (another JAK inhibitor) in rheumatoid arthritis patients 50 years and older with at least one additional cardiovascular risk factor, a higher rate of major adverse cardiovascular events (MACE), defined as cardiovascular death, non-fatal myocardial infarction (MI) and non-fatal stroke, was observed with tofacitinib compared to TNF inhibitors.

Therefore, in patients 65 years of age and older, patients who are current or past long-time smokers, and patients with history of atherosclerotic cardiovascular disease or other cardiovascular risk factors, abrocitinib should only be used if no suitable treatment alternatives are available.

### **Malignancy (excluding non-melanoma skin cancer [NMSC])**

Lymphoma and other malignancies have been reported in patients receiving JAK inhibitors, including abrocitinib.

In a large randomized active controlled study of tofacitinib (another JAK inhibitor) in rheumatoid arthritis patients 50 years and older with at least one additional cardiovascular risk factor, a higher rate of malignancies, particularly lung cancer, lymphoma and non-melanoma skin cancer (NMSC) was observed with tofacitinib compared to TNF inhibitors.

A higher rate of malignancies (excluding non-melanoma skin cancer, NMSC) was observed with abrocitinib 200 mg compared to abrocitinib 100 mg.

In patients 65 years of age and older, patients who are current or past long-time smokers, or with other malignancy risk factors (e.g. current malignancy or history of malignancy), abrocitinib should only be used if no suitable treatment alternatives are available.

### **Non-melanoma skin cancer**

NMSCs have been reported in patients receiving abrocitinib. Periodic skin examination is recommended for all patients, particularly those who are at increased risk for skin cancer.

## **Haematologic abnormalities**

Confirmed ALC  $<0.5 \times 10^3/\text{mm}^3$  and platelet count  $<50 \times 10^3/\text{mm}^3$  were observed in less than 0.5% of patients in clinical studies (see section 4.8). Treatment with abrocitinib should not be initiated in patients with a platelet count  $<150 \times 10^3/\text{mm}^3$ , an ALC  $<0.5 \times 10^3/\text{mm}^3$ , an ANC  $<1.2 \times 10^3/\text{mm}^3$  or who have a haemoglobin value  $<10 \text{ g/dL}$  (see section 4.2). Complete blood count should be monitored 4 weeks after initiation of therapy and thereafter according to routine patient management (see Table 1).

## **Lipids**

Dose-dependent increases in blood lipid parameters were reported in patients treated with abrocitinib compared to placebo (see section 4.8). Lipid parameters should be assessed approximately 4 weeks following initiation of therapy and thereafter according to the patient's risk for cardiovascular disease (see Table 1). The effect of these lipid parameter elevations on cardiovascular morbidity and mortality has not been determined. Patients with abnormal lipid parameters should be further monitored and managed according to clinical guidelines, due to the known cardiovascular risks associated with hyperlipidaemia.

## **Elderly**

The safety profile observed in elderly patients was similar to that of the adult population with the following exceptions: a higher proportion of patients 65 years of age and older discontinued from clinical studies and were more likely to have serious adverse reactions compared to younger patients; patients 65 years and older were more likely to develop low platelet and ALC values; the incidence rate of herpes zoster in patients 65 years of age and older was higher than that of younger patients (see section 4.8). There are limited data in patients above 75 years of age.

## **Use in patients 65 years of age and older**

Considering the increased risk of MACE, malignancies, serious infections, and all-cause mortality in patients 65 years of age and older, as observed in a large randomised study of tofacitinib (another JAK inhibitor), abrocitinib should only be used in these patients if no suitable treatment alternatives are available.

## **Immunosuppressive conditions or medicinal products**

Patients with immunodeficiency disorders or a first-degree relative with a hereditary immunodeficiency were excluded from clinical studies and no information on these patients is available.

Combination with biologic immunomodulators, potent immunosuppressants such as ciclosporin or other Janus kinase (JAK) inhibitors has not been studied. Their concomitant use with abrocitinib is not recommended as a risk of additive immunosuppression cannot be excluded.

## **Excipients**

### ***Lactose monohydrate***

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose galactose malabsorption should not take this medicinal product.

### ***Sodium***

This medicinal product contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

## **4.5. Interaction with Other Medicinal Products and Other Forms of Interaction**

### **Potential for other medicines to affect pharmacokinetics of abrocitinib**

Abrocitinib is metabolised predominantly by CYP2C19 and CYP2C9 enzymes, and to a lesser extent by CYP3A4 and CYP2B6 enzymes, and its active metabolites are renally excreted and are substrates of the organic anion transporter 3 (OAT3). Therefore, exposures of abrocitinib and/or its active metabolites may be affected by medicinal products that inhibit or induce these enzymes and transporter. Dose adjustments, as appropriate, are outlined in section 4.2.

### ***Co-administration with CYP2C19/CYP2C9 inhibitors***

When 100 mg abrocitinib was administered concomitantly with fluvoxamine (a strong CYP2C19 and moderate CYP3A inhibitor) or fluconazole (a strong CYP2C19, moderate CYP2C9 and

CYP3A inhibitor), the extent of exposure of abrocitinib active moiety (see section 5.2) increased by 91% and 155%, respectively, compared with administration alone (see section 4.2).

### ***Co-administration with CYP2C19/CYP2C9 inducers***

Administration of 200 mg abrocitinib after multiple doses with rifampicin, a strong inducer of CYP enzymes, resulted in reduction of abrocitinib active moiety exposures by approximately 56% (see section 4.2).

### ***Co-administration with OAT3 inhibitors***

When abrocitinib 200 mg was administered concomitantly with probenecid, an OAT3 inhibitor, abrocitinib active moiety exposures increased by approximately 66%. This is not clinically significant, and a dose adjustment is not needed.

### ***Co-administration with products which increase gastric pH***

When abrocitinib 200 mg was administered concomitantly with famotidine 40 mg, an H<sub>2</sub>-receptor antagonist, abrocitinib active moiety exposures decreased by approximately 35%. The effect of elevating gastric pH with antacids, or proton pump inhibitors (omeprazole) on the pharmacokinetics of abrocitinib has not been studied and may be similar to that seen with famotidine. The higher 200 mg daily dose should be considered for patients treated concomitantly with products which increase gastric pH, as they may reduce the efficacy of abrocitinib.

### **Potential for abrocitinib to affect pharmacokinetics of other medicinal products**

No clinically significant effects of abrocitinib were observed in interaction studies with oral contraceptives (e.g. ethinyl oestradiol/levonorgestrel).

*In vitro*, abrocitinib is an inhibitor of P glycoprotein (P-gp). Co-administration of dabigatran etexilate (a P-gp substrate), with a single dose of abrocitinib 200 mg increased dabigatran AUC<sub>inf</sub> and C<sub>max</sub> by approximately 53% and 40%, respectively, compared with administration alone. Caution should be exercised for concomitant use of abrocitinib with dabigatran. The effect of abrocitinib on the pharmacokinetics of other P-gp substrates has not been evaluated. Caution should be exercised as the levels of P-gp substrates with a narrow therapeutic index, such as

digoxin, may increase.

*In vitro*, abrocitinib is an inhibitor of CYP2C19 enzyme. Co-administration of abrocitinib 200 mg once daily with omeprazole 10 mg single dose increased the  $AUC_{inf}$  and  $C_{max}$  of omeprazole by approximately 189% and 134%, respectively, indicating that abrocitinib is a moderate inhibitor of CYP2C19 enzyme. Caution should be exercised when using abrocitinib concomitantly with narrow therapeutic index medicines that are primarily metabolised by CYP2C19 enzyme (e.g., S-mephenytoin and clopidogrel). Dose adjustment may be required for other medicines primarily metabolised by CYP2C19 enzyme in accordance with their product information (e.g., citalopram, clobazam, escitalopram and selumetinib).

Co-administration of abrocitinib 200 mg once daily with caffeine 100 mg single dose increased the  $AUC_{inf}$  of caffeine by 40% with lack of effect on  $C_{max}$ , suggesting that abrocitinib is a mild inhibitor of CYP1A2 enzyme. No general dose adjustment can be recommended.

#### **4.6. Fertility, Pregnancy and Lactation**

##### **Women of childbearing potential**

Women of reproductive potential should be advised to use effective contraception during treatment and for 1 month following the final dose of CIBINQO. Pregnancy planning and prevention for females of reproductive potential should be encouraged.

##### **Pregnancy**

There are no or limited amount of data on the use of abrocitinib in pregnant women. Studies in animals have shown reproductive toxicity. Abrocitinib has been shown to cause embryo-foetal lethality in pregnant rats and rabbits, skeletal variations in the foetuses of pregnant rats and rabbits, and to affect parturition and peri/postnatal development in rats (see section 5.3). CIBINQO is contraindicated during pregnancy (see section 4.3).

##### **Breast-feeding**

There are no data on the presence of abrocitinib in human milk, the effects on the breast-fed infant, or the effects on milk production. Abrocitinib was secreted in milk of lactating rats. A risk to

newborns/infants cannot be excluded and CIBINQO is contraindicated during breast-feeding (see section 4.3).

## **Fertility**

Based on the findings in rats, oral administration of CIBINQO may result in temporary reduced fertility in females of reproductive potential. The effects on female rat fertility were reversible 1 month after cessation of abrocitinib oral administration (see section 5.3).

### **4.7. Effects on Ability to Drive and Use Machines**

CIBINQO has no or negligible influence on the ability to drive and use machines.

### **4.8. Undesirable Effects**

#### **Summary of the safety profile**

The most commonly reported adverse reactions are nausea (15.1%), headache (7.9%), acne (4.8%), herpes simplex (4.2%), blood creatine phosphokinase increased (3.8%), vomiting (3.5%), dizziness (3.4%) and abdominal pain upper (2.2%). The most frequent serious adverse reactions are infections (0.3%) (see section 4.4).

#### **Tabulated list of adverse reactions**

A total of 3848 patients were treated with abrocitinib in clinical studies in atopic dermatitis. Among them 3050 patients (representing 5166 patient-years of exposure) were integrated for safety analysis. The integrated safety analysis included 1997 patients receiving a constant dose of abrocitinib 200 mg and 1053 patients receiving a constant dose of 100 mg. There were 2013 patients with at least 48 weeks of exposure. Five placebo-controlled studies were integrated (703 patients on 100 mg once daily, 684 patients on 200 mg once daily and 438 patients on placebo) to evaluate the safety of abrocitinib in comparison to placebo for up to 16 weeks.

Listed in Table 2 are adverse reactions observed in atopic dermatitis clinical studies presented by system organ class and frequency, using the following categories: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1000$  to  $< 1/100$ ); rare ( $\geq 1/10000$  to  $< 1/1000$ ); very rare

(<1/10000). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

**Table 2. Adverse reactions**

System organ class	Very common	Common	Uncommon
Infections and infestations		Herpes simplex <sup>a</sup> Herpes zoster <sup>b</sup>	Pneumonia
Blood and lymphatic system disorders			Thrombocytopenia Lymphopenia
Metabolism and nutrition disorders			Hyperlipidaemia <sup>c</sup>
Nervous system disorders		Headache Dizziness	
Vascular disorders			Venous thromboembolism <sup>d</sup>
Gastrointestinal disorders	Nausea	Vomiting Abdominal pain upper	
Skin and subcutaneous tissue disorders		Acne	
Investigations		Creatine phosphokinase increased >5 × ULN <sup>e</sup>	

a. Herpes simplex includes oral herpes, ophthalmic herpes simplex, genital herpes, and herpes dermatitis.

b. Herpes zoster includes ophthalmic herpes zoster.

c. Hyperlipidaemia includes dyslipidaemia and hypercholesterolaemia.

d. Venous thromboembolism includes pulmonary embolism and deep vein thrombosis.

e. Includes changes detected during laboratory monitoring (see text below).

## Description of selected adverse reactions

### ***Infections***

In placebo-controlled studies, for up to 16 weeks, infections have been reported in 27.4% of patients treated with placebo and in 34.9% and 34.8% of patients treated with abrocitinib 100 mg



and 200 mg, respectively. Most infections were mild or moderate. The percentage of patients reporting infection-related adverse reactions in the 200 mg and 100 mg groups compared to placebo were: herpes simplex (4.2% and 2.8% versus 1.4%), herpes zoster (1.2% and 0.6% versus 0%), pneumonia (0.1% and 0.1% versus 0%). Herpes simplex was more frequent in patients with a history of herpes simplex or eczema herpeticum. Most of the herpes zoster events involved a single dermatome and were non-serious. Most opportunistic infections were cases of herpes zoster (0.70 per 100 patient-years in the abrocitinib 100 mg group and 0.96 per 100 patient-years in the abrocitinib 200 mg group), most of which were non-serious multidermatomal cutaneous infections. Among all patients treated in clinical studies with consistent dosing regimens of either abrocitinib 100 mg or 200 mg, including the long-term extension study, the incidence rate of herpes zoster in patients treated with abrocitinib 200 mg (4.36 per 100 patient-years) was higher than that of patients treated with 100 mg (2.61 per 100 patient-years). Incidence rates for herpes zoster were also higher for patients 65 years of age and older (HR 1.76), patients with a medical history of herpes zoster (HR 3.41), patients with severe atopic dermatitis at baseline (HR 1.17), and a confirmed ALC  $<1.0 \times 10^3/\text{mm}^3$  prior to the event of herpes zoster (HR 2.18) (see section 4.4).

In placebo-controlled studies, for up to 16 weeks, the rate of serious infections was 1.81 per 100 patient-years in patients treated with placebo, 3.32 per 100 patient-years in patients treated with 100 mg, and 1.12 per 100 patient-years in patients treated with 200 mg. Among all patients treated in clinical studies with consistent dosing regimens of either abrocitinib 100 mg or 200 mg, including the long-term extension study, the rate of serious infections was 2.20 per 100 patient-years treated with 100 mg and 2.46 per 100 patient-years treated with 200 mg. The most commonly reported serious infections were herpes simplex, herpes zoster, and pneumonia (see section 4.4).

### ***Venous thromboembolism***

Among all patients treated in clinical studies with consistent dosing regimens of either abrocitinib 100 mg or 200 mg, including the long-term extension study, the rate of PE was 0.21 per 100 patient-years for 200 mg and 0.05 per 100 patient-years for 100 mg. The rate of DVT was 0.06 per 100 patient-years in the 200 mg group and 0.05 per 100 patient-years in the 100 mg group (see section 4.4).

### ***Thrombocytopenia***

In placebo-controlled studies, for up to 16 weeks, treatment was associated with a dose-related decrease in platelet count. Maximum effects on platelets were observed within 4 weeks, after which the platelet count returned towards baseline despite continued therapy. Confirmed platelet counts of  $<50 \times 10^3/\text{mm}^3$  were reported in 0.1% of patients exposed to 200 mg, and in 0 patients treated with 100 mg or placebo. Among all patients treated with clinical studies with consistent dosing regimens of either abrocitinib 100 mg or 200 mg, including the long-term extension study, the rate of confirmed platelet counts of  $<50 \times 10^3/\text{mm}^3$  was 0.15 per 100 patients-years for 200 mg and 0 per 100 patient years for 100 mg, most occurring at Week 4. Patients 65 years of age and older had a higher rate of platelet counts  $<75 \times 10^3/\text{mm}^3$  (see section 4.4).

### ***Lymphopenia***

In placebo-controlled studies, for up to 16 weeks, confirmed ALC  $<0.5 \times 10^3/\text{mm}^3$  occurred in 0.3% of patients treated with 200 mg and 0% of patients treated with 100 mg or placebo. Both cases occurred in the first 4 weeks of exposure. Among all patients treated in clinical studies with consistent dosing regimens of either abrocitinib 100 mg or 200 mg, including the long-term extension, the rate of confirmed ALC  $<0.5 \times 10^3/\text{mm}^3$  was 0.34 per 100 patient-years for 200 mg and 0 per 100 patient-years for 100 mg, the highest rate was observed in patients 65 years of age and older (see section 4.4).

### ***Lipid elevations***

In placebo-controlled studies, for up to 16 weeks, there was a dose-related increase in low-density lipoprotein cholesterol (LDL-c), total cholesterol, and high-density lipoprotein cholesterol (HDL-c) relative to placebo at Week 4 which remained elevated through the final visit in the treatment period. There was no meaningful change in the LDL/HDL ratio in patients treated with abrocitinib relative to patients treated with placebo. Events related to hyperlipidaemia occurred in 0.4% of patients exposed to abrocitinib 100 mg, 0.6% of patients exposed to 200 mg and 0% of patients exposed to placebo (see section 4.4).

### ***Creatine phosphokinase elevations (CPK)***

In placebo-controlled studies, for up to 16 weeks, significant increases in CPK values ( $>5 \times \text{ULN}$ ) occurred in 1.8% of patients treated with placebo, 1.8% of patients treated with 100 mg and 3.8%

of patients treated with 200 mg of abrocitinib, respectively. Most elevations were transient, and none led to discontinuation.

### **Nausea**

In placebo-controlled studies, for up to 16 weeks, nausea was reported in 1.8% of patients treated with placebo and in 6.3% and 15.1% of patients treated with 100 mg and 200 mg, respectively. Discontinuation due to nausea occurred in 0.4% of patients treated with abrocitinib. Among patients with nausea, 63.5% of patients had onset of nausea in the first week of therapy. The median duration of nausea was 15 days. Most of the cases were mild to moderate in severity.

### **Paediatric population**

A total of 635 adolescent patients (12 to less than 18 years of age) were treated with abrocitinib in clinical studies in atopic dermatitis representing 1326.1 patient-years of exposure. The safety profile observed in adolescents in atopic dermatitis clinical studies was similar to that of the adult population.

### **4.9. Overdose**

CIBINQO was administered in clinical studies up to a single oral dose of 800 mg and 400 mg daily for 28 days. Adverse reactions were comparable to those seen at lower doses and no specific toxicities were identified. In case of an overdose, it is recommended that the patient be monitored for signs and symptoms of adverse reactions (see section 4.8). Treatment should be symptomatic and supportive. There is no specific antidote for overdose with this medicinal product.

Pharmacokinetics data up to and including a single oral dose of 800 mg in healthy adult volunteers indicate that more than 90% of the administered dose is expected to be eliminated within 48 hours.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1. Pharmacodynamic Properties**

Pharmacotherapeutic group: Other dermatological preparations, agents for dermatitis, excluding corticosteroids; ATC code: D11AH08

### **Mechanism of action**

Abrocitinib is a Janus kinase (JAK) 1 inhibitor. JAKs are intracellular enzymes which transmit signals arising from cytokine or growth factor-receptor interactions on the cellular membrane to influence cellular processes of haematopoiesis and immune cell function. JAKs phosphorylate and activate Signal Transducers and Activators of Transcription (STATs) which modulate intracellular activity including gene expression. Inhibition of JAK1 modulates the signalling pathways by preventing the phosphorylation and activation of STATs.

In biochemical assays, abrocitinib has selectivity for JAK1 over the other 3 JAK isoforms JAK2 (28-fold), JAK3 (>340-fold) and tyrosine kinase 2 (TYK2, 43-fold). In cellular settings, it preferentially inhibits cytokine-induced STAT phosphorylation by signalling pairs involving JAK1, and spares signalling by JAK2/JAK2, or JAK2/TYK2 pairs. The relevance of selective enzymatic inhibition of specific JAK enzymes to clinical effect is not currently known.

### **Pharmacodynamic effects**

#### ***Clinical biomarkers***

Treatment with abrocitinib was associated with dose-dependent reduction in serum biomarkers of inflammation in atopic dermatitis [interleukin-31 (IL-31), interleukin-22 (IL-22), eosinophil count, and thymus and activation-regulated chemokine (TARC)], JAK1 signalling [natural killer (NK) cell count and interferon gamma-induced protein 10 (IP-10)] or both [high sensitivity C-reactive protein (hsCRP)]. These changes were reversible after treatment discontinuation.

Mean absolute lymphocyte count increased by 2 weeks after starting treatment with abrocitinib and returned to baseline by Month 9 of treatment. Most patients maintained an ALC within the reference range. Treatment with abrocitinib was associated with a dose-related increase in B cell counts and a dose-related decrease in NK cell counts. The clinical significance of these changes in B cell and NK cell counts is unknown.

### ***Cardiac electrophysiology***

The effect of abrocitinib on the QTc interval was examined in subjects who received a single suprathreshold dose of abrocitinib 600 mg in a placebo- and positive-controlled thorough QT study. A concentration-dependent QTc prolonging effect of abrocitinib was seen; the mean (90% confidence interval) for the increase in QTc interval was 6.0 (4.52, 7.49) msec, indicating the lack of a clinically relevant effect of abrocitinib on QTc interval at the dose tested.

### **Clinical efficacy and safety**

The efficacy and safety of abrocitinib as monotherapy and in combination with background medicated topical therapies over 12-16 weeks were evaluated in 1616 patients in 3 pivotal Phase 3 randomised, double-blind, placebo-controlled studies (MONO-1, MONO-2, and COMPARE). In addition, the efficacy and safety of abrocitinib in monotherapy over 52 weeks (with the option of rescue treatment in flaring patients) was evaluated in 1233 patients in a Phase 3 induction, randomised withdrawal, double-blind, placebo-controlled study (REGIMEN). The patients in these 4 studies were 12 years of age and older with moderate-to-severe atopic dermatitis as defined by Investigator's Global Assessment (IGA) score  $\geq 3$ , Eczema Area and Severity Index (EASI) score  $\geq 16$ , body surface area (BSA) involvement  $\geq 10\%$ , and Peak Pruritus Numerical Rating Scale (PP-NRS)  $\geq 4$  at baseline prior to randomisation. Patients who had a prior inadequate response or for whom topical treatments were medically inadvisable, or who had received systemic therapies were eligible for inclusion. All patients who completed the parent studies were eligible to enroll into the long-term extension study EXTEND.

### ***Baseline characteristics***

In the placebo-controlled studies (MONO-1, MONO-2, COMPARE) and the open-label induction, randomised withdrawal study (REGIMEN), across all treatment groups 41.4% to 51.1% were female, 59.3% to 77.8% were Caucasian, 15.0% to 33.0% were Asian and 4.1% to 8.3% were Black, and the mean age was 32.1 to 37.7 years. A total of 134 patients 65 years of age and older were enrolled in these studies. In these studies, 32.2% to 40.8% had a baseline IGA of 4 (severe atopic dermatitis), and 41.4% to 59.5% of patients had received prior systemic treatment for atopic dermatitis. The baseline mean EASI score ranged from 28.5 to 30.9, the baseline PP-NRS ranged from 7.0 to 7.3 and the baseline Dermatology Life Quality Index (DLQI) ranged from 14.4 to 16.0.

## Clinical response

### 12-week monotherapy (MONO-1, MONO-2) and 16-week combination therapy (COMPARE) studies

A significantly larger proportion of patients achieved both primary endpoints IGA 0 or 1 and/or EASI-75 with 100 mg or 200 mg once daily abrocitinib compared with placebo at Week 12 or Week 16 (see Table 3 and Table 4).

A significantly greater proportion of patients achieved at least a PP-NRS 4-point improvement with 100 mg or 200 mg once daily abrocitinib compared with placebo. This improvement was observed as early as Week 2 and persisted through Week 12 (Figure 1).

In the COMPARE study, superiority of abrocitinib 200 mg compared with dupilumab at Week 2 was demonstrated for the proportion of patients achieving at least a PP-NRS 4-point improvement with significantly higher itch responses seen as early as Day 4 after the first dose.

Treatment effects in subgroups (e.g., weight, age, sex, race and prior systemic immunosuppressant treatment) in MONO-1, MONO-2 and COMPARE were consistent with the results in the overall study population.

**Table 3. Efficacy results of abrocitinib in monotherapy at Week 12**

	MONO-1 <sup>d</sup>			MONO-2 <sup>d</sup>		
	Week 12			Week 12		
	Abrocitinib monotherapy		PBO	Abrocitinib monotherapy		PBO
	200 mg QD N=154	100 mg QD N=156		200 mg QD N=155	100 mg QD N=158	
	<b>% Responders (95% CI)</b>					
IGA 0 or 1 <sup>a</sup>	43.8 <sup>e</sup> (35.9, 51.7)	23.7 <sup>e</sup> (17.0, 30.4)	7.9 (1.8, 14.0)	38.1 <sup>e</sup> (30.4, 45.7)	28.4 <sup>e</sup> (21.3, 35.5)	9.1 (2.7, 15.5)
EASI-75 <sup>b</sup>	62.7 <sup>e</sup> (55.1, 70.4)	39.7 <sup>e</sup> (32.1, 47.4)	11.8 (4.6, 19.1)	61.0 <sup>e</sup> (53.3, 68.7)	44.5 <sup>e</sup> (36.7, 52.3)	10.4 (3.6, 17.2)
PP-NRS4 <sup>c</sup>	57.2 <sup>e</sup> (48.8, 65.6)	37.7 <sup>e</sup> (29.2, 46.3)	15.3 (6.6, 24.0)	55.3 <sup>e</sup> (47.2, 63.5)	45.2 <sup>e</sup> (37.1, 53.3)	11.5 (4.1, 19.0)

**Table 3. Efficacy results of abrocitinib in monotherapy at Week 12**

	MONO-1 <sup>d</sup>			MONO-2 <sup>d</sup>		
	Week 12			Week 12		
	Abrocitinib monotherapy		PBO	Abrocitinib monotherapy		PBO
	200 mg QD N=154	100 mg QD N=156		200 mg QD N=155	100 mg QD N=158	

Abbreviations: CI=confidence interval; EASI=Eczema Area and Severity Index; IGA=Investigator Global Assessment; N=number of patients randomised; PBO=placebo; PP-NRS=Peak Pruritus Numerical Rating Scale; QD=once daily.

- IGA responders were patients with IGA score of clear (0) or almost clear (1) (on a 5-point scale) and a reduction from baseline of  $\geq 2$  points.
- EASI-75 responders were patients with  $\geq 75\%$  improvement in EASI from baseline.
- PP-NRS4 responders were patients with  $\geq 4$ -point improvement in PP-NRS from baseline.
- Abrocitinib used as monotherapy.
- Statistically significant with adjustment for multiplicity versus placebo.

**Table 4. Efficacy results of abrocitinib in combination with topical therapy at Week 12 and Week 16**

	COMPARE <sup>d</sup>							
	Week 12				Week 16			
	Abrocitinib + topicals		PBO + topicals N=131	DUP + topicals N=243	Abrocitinib + topicals		PBO + topicals N=131	DUP + topicals N=243
	200 mg QD N=226	100 mg QD N=238			200 mg QD N=226	100 mg QD N=238		
	<b>% Responders (95% CI)</b>							
IGA 0 or 1 <sup>a</sup>	48.4 <sup>e</sup> (41.8, 55.0)	36.6 <sup>e</sup> (30.4, 42.8)	14.0 (8.0, 19.9)	36.5 (30.4, 42.6)	47.5 <sup>e</sup> (40.9, 54.1)	34.8 <sup>e</sup> (28.6, 40.9)	12.9 (7.0, 18.8)	38.8 (32.5, 45.1)
EASI-75 <sup>b</sup>	70.3 <sup>e</sup> (64.3, 76.4)	58.7 <sup>e</sup> (52.4, 65.0)	27.1 (19.5, 34.8)	58.1 (51.9, 64.3)	71.0 <sup>e</sup> (65.1, 77.0)	60.3 <sup>e</sup> (53.9, 66.6)	30.6 (22.5, 38.8)	65.5 (59.4, 71.6)
PP-NRS4 <sup>c</sup>	63.1 (56.7, 69.6)	47.5 (40.9, 54.1)	28.9 (20.8, 37.0)	54.5 (47.9, 61.0)	62.8 (55.6, 70.0)	47.0 (39.5, 54.6)	28.7 (19.6, 37.9)	57.1 (50.1, 64.2)

Abbreviations: CI=confidence interval; DUP=Dupilumab; EASI=Eczema Area and Severity Index; IGA=Investigator Global Assessment; N=number of patients randomised; PBO=placebo; PP-NRS=Peak Pruritus Numerical Rating Scale; QD=once daily.

**Table 4. Efficacy results of abrocitinib in combination with topical therapy at Week 12 and Week 16**


	COMPARE <sup>d</sup>							
	Week 12				Week 16			
	Abrocitinib + topicals		PBO + topicals	DUP + topicals	Abrocitinib + topicals		PBO + topicals	DUP + topicals
	200 mg QD N=226	100 mg QD N=238	N=131	N=243	200 mg QD N=226	100 mg QD N=238	N=131	N=243

- a. IGA responders were patients with IGA score of clear (0) or almost clear (1) (on a 5-point scale) and a reduction from baseline of  $\geq 2$  points.
- b. EASI-75 responders were patients with  $\geq 75\%$  improvement in EASI from baseline.
- c. PP-NRS4 responders were patients with  $\geq 4$ -point improvement in PP-NRS from baseline.
- d. Abrocitinib used in combination with topical therapy.
- e. Statistically significant with adjustment for multiplicity versus placebo.

The proportion of patients who achieved PP-NRS4 over time in studies MONO-1, MONO-2 and COMPARE are shown in Figure 1.



**Figure 1. Proportion of patients who achieved PP-NRS4 over time in MONO-1, MONO-2 and COMPARE**



Abbreviations: PP-NRS=Peak Pruritus Numerical Rating Scale; QD=once daily; Q2W=every 2 weeks.

PP-NRS4 responders were patients with  $\geq 4$ -point improvement in PP-NRS from baseline.

- a. Abrocitinib used as monotherapy.
- b. Abrocitinib used in combination with medicated topical therapy.
- \* Statistically significant with adjustment for multiplicity versus placebo.
- \*\* Statistically significant with adjustment for multiplicity versus dupilumab.

Health-related outcomes

In both monotherapy studies (MONO-1 and MONO-2) and in the combination therapy study (COMPARE), abrocitinib significantly improved patient-reported outcomes, including itch, sleep (SCORAD Sleep VAS), AD symptoms (POEM), quality of life (DLQI) and symptoms of anxiety and depression (HADS) that were uncorrected for multiplicity, at 12 weeks compared to placebo (see Table 5).

**Table 5. Patient-reported outcomes results of abrocitinib monotherapy and in combination with topical therapy at Week 12**

	Monotherapy						Combination therapy		
	MONO-1			MONO-2			COMPARE		
	200 mg QD	100 mg QD	PBO	200 mg QD	100 mg QD	PBO	200 mg QD + topicals	100 mg QD + topicals	PBO + topicals
<b>N</b>	<b>154</b>	<b>156</b>	<b>77</b>	<b>155</b>	<b>158</b>	<b>78</b>	<b>226</b>	<b>238</b>	<b>131</b>
SCORAD Sleep VAS, change from baseline (95% CI)	-3.7* (-4.2, -3.3)	-2.9* (-3.4, -2.5)	-1.6 (-2.2, -1.0)	-3.8* (-4.2, -3.4)	-3.0* (-3.4, -2.6)	-2.1 (-2.7, -1.5)	-4.6* (-4.9, -4.3)	-3.7* (-4.0, -3.4)	-2.4 (-2.8, -2.0)
DLQI $\geq$ 4-point improvement, % responders	72.6%*	67.2%*	43.6%	78.1%*	73.3%*	32.3%	86.4%*	74.7%*	56.5%
POEM, change from baseline (95% CI)	-10.6* (-11.8, -9.4)	-6.8* (-8.0, -5.6)	-3.7 (-5.5, -1.9)	-11.0* (-12.1, -9.8)	-8.7* (-9.9, -7.5)	-3.6 (-5.3, -1.9)	-12.6* (-13.6, -11.7)	-9.6* (-10.5, -8.6)	-5.1 (-6.3, -3.9)
HADS Anxiety, change from baseline (95% CI)	-2.1* (-2.5, -1.6)	-1.6 (-2.0, -1.1)	-1.0 (-1.7, -0.4)	-1.7* (-2.2, -1.2)	-1.6* (-2.1, -1.1)	-0.6 (-1.3, -0.2)	-1.6* (-2.0, -1.2)	-1.2* (-1.5, -0.8)	-0.4 (-0.9, 0.1)
HADS Depression, change from baseline (95% CI)	-1.8* (-2.2, -1.4)	-1.4* (-1.8, -0.9)	-0.2 (-0.8, 0.4)	-1.4* (-1.8, -1.0)	-1.0* (-1.5, -0.6)	0.3 (-0.3, 0.9)	-1.6* (-1.9, -1.2)	-1.3* (-1.6, -0.9)	-0.3 (-0.7, 0.2)

CI=confidence interval; DLQI=Dermatology Life Quality Index; HADS=Hospital Anxiety and Depression Scale; N=number of patients randomised; PBO=placebo; POEM=Patient-Oriented Eczema Measure; QD=once daily; SCORAD=SCORing for AD; VAS=visual analog scale.

\*Statistically significant without adjusting for multiplicity


Open-label induction, randomised withdrawal study (REGIMEN)

A total of 1233 patients received open-label abrocitinib 200 mg once daily in the 12-week run-in

phase. Among these patients, 798 patients (64.7%) met responder criteria (defined as achieving IGA [0 or 1] response and EASI-75) and were randomised to placebo (267 patients), abrocitinib 100 mg once daily (265 patients) or abrocitinib 200 mg once daily (266 patients).

Continuous treatment (200 mg continuous) and induction-maintenance treatment (200 mg for 12 weeks followed by 100 mg) prevented flare with 81.1% and 57.4% probability, respectively, versus 19.1% among patients who withdrew treatment (randomised to placebo) after 12 weeks of induction. Three hundred fifty-one (351) patients including 16.2% of 200 mg, 39.2% of 100 mg and 76.4% of placebo patients received rescue medication of 200 mg abrocitinib in combination with topical therapy.

**Figure 2. Time to protocol-defined flare**



Abrocitinib used as monotherapy.

Protocol-defined flare=A loss of at least 50% of the EASI response at Week 12 and an IGA score of 2 or higher.

Multiplicity-controlled p <0.0001 200 mg versus placebo; 100 mg versus placebo; 200 mg versus 100 mg.

Long-term efficacy

Eligible patients who completed the full treatment period of a qualifying parent study (e.g., MONO-1, MONO-2, COMPARE, REGIMEN) were considered for enrolment in the long-term extension study EXTEND. In EXTEND, patients received abrocitinib with or without background medicated topical therapy. Patients who were previously randomised to medicinal product 100 mg

or 200 mg once daily in parent studies continued the same dose in EXTEND as in the parent study. In EXTEND, patients received double-blind treatment until the parent study was completed, after which patients received single-blind treatment (treatment assignment disclosed to the investigators but not to the patients).

Among patients who achieved response after 12 weeks of treatment and entered EXTEND, the majority of patients maintained their response at Week 96 of cumulative treatment for both doses of abrocitinib [64% and 72% for IGA (0 or 1) response, 87% and 90% for EASI-75, and 75% and 80% for PP-NRS4 with 100 mg once daily and 200 mg once daily, respectively].

Among patients who did not achieve response after 12 weeks of treatment and entered EXTEND, a proportion of patients achieved late-onset response by Week 24 (from baseline) of continued treatment with abrocitinib [25% and 29% for IGA (0 or 1) response, and 50% and 57% for EASI-75 with 100 mg once daily and 200 mg once daily, respectively]. Patients who achieved partial response at Week 12 were more likely than those with no response at Week 12 to achieve treatment benefit at Week 24.

Patients who received dupilumab in the COMPARE study and subsequently entered EXTEND were randomised to either 100 mg or 200 mg of abrocitinib once daily upon entering EXTEND. Among non-responders to dupilumab, a substantial proportion of patients achieved response 12 weeks after switching to abrocitinib [34% and 47% for IGA (0 or 1) response, and 68% and 80% for EASI-75 with 100 mg once daily or 200 mg once daily, respectively].

### **Paediatric population**

The efficacy and safety of 12 weeks of abrocitinib monotherapy were evaluated in 2 Phase 3 randomised, double-blind, placebo-controlled studies (MONO-1, MONO-2) which included 124 patients who were 12 to less than 18 years of age. The efficacy and safety of abrocitinib monotherapy over 52 weeks (with the option of rescue treatment for flaring patients) were also evaluated in an open-label induction, randomised withdrawal study (REGIMEN), which included 246 patients who were 12 to less than 18 years of age. In these studies, the results in the adolescent subgroup were consistent with the results in the overall study population.

The efficacy and safety of 12 weeks of abrocitinib in combination with background medicated topical therapy were evaluated in the Phase 3 randomised, double-blind, placebo-controlled study

TEEN. The study included 287 patients who were 12 to less than 18 years of age with moderate-to-severe atopic dermatitis as defined by IGA score  $\geq 3$ , EASI score  $\geq 16$ , BSA involvement  $\geq 10\%$ , and PP-NRS  $\geq 4$  at the baseline visit prior to randomisation. Patients who had a prior inadequate response or who had received systemic therapy, were eligible for inclusion.

### Baseline characteristics

In TEEN, across all treatment groups 49.1% were female, 56.1% were Caucasian, 33.0% were Asian and 6.0% were Black patients. The median age was 15 years and the proportion of patients with severe atopic dermatitis (IGA of 4) was 38.6%.

Results of 12-week abrocitinib treatment in adolescents in pooled MONO-1 and MONO-2, and the TEEN study are shown in Table 6.

**Table 6. Adolescent efficacy results at Week 12 in pooled MONO-1 and MONO-2, and in TEEN**

	Pooled MONO-1 and MONO-2			TEEN <sup>d</sup>		
	Abrocitinib 200 mg QD	Abrocitinib 100 mg QD	Placebo	Abrocitinib 200 mg QD	Abrocitinib 100 mg QD	Placebo
<b>IGA 0 or 1<sup>a</sup></b>						
N	48	50	23	93	89	94
%	31.3	22.0	8.7	46.2 <sup>e</sup>	41.6 <sup>e</sup>	24.5
95% CI	(18.1, 44.4)	(10.5, 33.5)	(0.0, 20.2)	(36.1, 56.4)	(31.3, 51.8)	(15.8, 33.2)
<b>EASI-75<sup>b</sup></b>						
N	48	50	23	93	89	94
%	56.3	44.0	8.7	72.0 <sup>e</sup>	68.5 <sup>e</sup>	41.5
95% CI	(42.2, 70.3)	(30.2, 57.8)	(0.0, 20.2)	(62.9, 81.2)	(58.9, 78.2)	(31.5, 51.4)
<b>PP-NRS4<sup>c</sup></b>						
N	36	42	22	74	76	84
%	61.1	28.6	9.1	55.4 <sup>e</sup>	52.6	29.8
95% CI	(45.2, 77.0)	(14.9, 42.2)	(0.0, 21.1)	(44.1, 66.7)	(41.4, 63.9)	(20.0, 39.5)

Abbreviations: CI=confidence interval; EASI=Eczema Area and Severity Index; IGA=Investigator Global Assessment; N=number of evaluable patients; PP-NRS=Peak Pruritus Numerical Rating Scale; QD=once daily.

a. IGA responders were patients with IGA score of clear (0) or almost clear (1) (on a 5 point scale) and a reduction from

baseline of  $\geq 2$  points.

- b. EASI-75 responders were patients with  $\geq 75\%$  improvement in EASI from baseline.
- c. PP-NRS4 responders were patients with  $\geq 4$ -point improvement in PP-NRS from baseline.
- d. Abrocitinib used in combination with medicated topical therapy.
- e. Statistically significant with adjustment for multiplicity versus placebo.

Among adolescent patients who achieved response after 12 weeks of treatment and entered long-term extension study EXTEND, the majority of patients maintained their response at Week 96 of cumulative treatment for both doses of abrocitinib [62% and 78% for IGA (0 or 1) response, 89% and 93% for EASI-75, and 77% and 76% for PP-NRS4 with 100 mg and 200 mg once daily, respectively].

Among adolescent patients who did not achieve response after 12 weeks of treatment and entered EXTEND, a proportion of patients achieved late-onset response by Week 24 (from baseline) of continued treatment with both doses of abrocitinib [34% and 28% for IGA (0 or 1) response, and 41% and 55% for EASI-75 with 100 mg and 200 mg once daily, respectively].

## 5.2. Pharmacokinetic Properties

### Absorption

Abrocitinib is well-absorbed with over 91% extent of oral absorption and absolute oral bioavailability of approximately 60%. The oral absorption of abrocitinib is rapid and peak plasma concentrations are reached within 1 hour. Steady-state plasma concentrations of abrocitinib are achieved within 48 hours after once daily administration. Both  $C_{\max}$  and AUC of abrocitinib increased dose proportionally up to 200 mg. Co-administration of abrocitinib with a high-fat meal had no clinically relevant effect on abrocitinib exposures (AUC and  $C_{\max}$  increased by approximately 26% and 29%, respectively, and  $T_{\max}$  was prolonged by 2 hours). In clinical studies, abrocitinib was administered without regard to food (see section 4.2).

### Distribution

After intravenous administration, the volume of distribution of abrocitinib is about 100 L. Approximately 64%, 37% and 29% of circulating abrocitinib and its active metabolites M1 and M2, respectively, are bound to plasma proteins. Abrocitinib and its active metabolites distribute equally

between red blood cells and plasma.

## **Biotransformation**

The *in vitro* metabolism of abrocitinib is mediated by multiple CYP enzymes, CYP2C19 (~53%), CYP2C9 (~30%), CYP3A4 (~11%) and CYP2B6 (~6%). In a human radiolabelled study, abrocitinib was the most prevalent circulating species, with mainly 3 polar mono-hydroxylated metabolites identified as M1 (3-hydroxypropyl), M2 (2-hydroxypropyl) and M4 (pyrrolidinone pyrimidine). At steady state, M2 and M4 are major metabolites and M1 is a minor metabolite. Of the 3 metabolites in circulation, M1 and M2 have similar JAK inhibitory profiles as abrocitinib, while M4 was pharmacologically inactive. The pharmacologic activity of abrocitinib is attributable to the unbound exposures of parent molecule (~60%) as well as M1 (~10%) and M2 (~30%) in systemic circulation. The sum of unbound exposures of abrocitinib, M1 and M2, each expressed in molar units and adjusted for relative potencies, is referred to as the abrocitinib active moiety.

No clinically significant effects of abrocitinib were observed in interaction studies with substrates of BCRP and OAT3 (e.g. rosuvastatin), MATE1/2K (e.g. metformin), CYP3A4 (e.g. midazolam), and CYP2B6 (e.g. efavirenz).

## **Elimination**

The elimination half-life of abrocitinib is about 5 hours. Abrocitinib is eliminated primarily by metabolic clearance mechanisms, with less than 1% of the dose excreted in urine as unchanged active substance. The metabolites of abrocitinib, M1, M2 and M4 are excreted predominantly in urine, and are substrates of OAT3 transporter.

## **Special populations**

### ***Body weight, gender, genotype, race, and age***

Body weight, gender, CYP2C19/2C9 genotype, race, and age did not have a clinically meaningful effect on abrocitinib exposure (see section 4.2).

### ***Adolescents ( $\geq 12$ to $< 18$ years)***

Based on population pharmacokinetic analysis, there was no clinically relevant difference in mean abrocitinib steady-state exposures in adolescent patients compared to adults at their typical body weights.

### ***Paediatric (<12 years)***

Interaction studies have been performed in adults only. The pharmacokinetics of abrocitinib in children under 12 years of age have not yet been established (see section 4.2).

### **Renal impairment**

In a renal impairment study, patients with severe (eGFR <30 mL/min) and moderate (eGFR 30 to <60 mL/min) renal impairment had approximately 191% and 110% increase in active moiety  $AUC_{inf}$ , respectively, compared to patients with normal renal function (eGFR  $\geq$ 90 mL/min) (see section 4.2). Pharmacokinetics of abrocitinib have not been determined in patients with mild renal impairment, however, based on the results observed in other groups, an increase of up to 70% in active moiety exposure is expected in patients with mild renal impairment (eGFR 60 to <90 mL/min). The increase of up to 70% is not clinically meaningful as the efficacy and safety of abrocitinib in atopic dermatitis patients with mild renal impairment (n=756) was comparable to the overall population in Phase 2 and 3 clinical studies. The eGFR in individual patients was estimated using Modification of Diet in Renal Disease (MDRD) formula.

Abrocitinib has not been studied in patients with ESRD on renal replacement therapy (see section 4.2). In Phase 3 clinical studies, abrocitinib was not evaluated in patients with atopic dermatitis with baseline creatinine clearance values less than 40 mL/min.

### **Hepatic impairment**

Patients with mild (Child Pugh A) and moderate (Child Pugh B) hepatic impairment had approximately 4% decrease and 15% increase in active moiety  $AUC_{inf}$ , respectively, compared to patients with normal hepatic function. These changes are not clinically significant, and no dose adjustment is required in patients with mild or moderate hepatic impairment (see section 4.2). In clinical studies, abrocitinib was not evaluated in patients with severe (Child Pugh C) hepatic impairment (see section 4.3), or in patients screened positive for active hepatitis B or hepatitis C (see section 4.4).



### 5.3. Preclinical Safety Data

#### General toxicity

Decreased lymphocyte counts and decreased size and/or lymphoid cellularity of organs/tissues of the immune and haematopoietic systems were observed in nonclinical studies and were attributed to the pharmacological properties (JAK inhibition) of abrocitinib.

In toxicity studies of up to 1 month of abrocitinib dosing in rats at an age comparable to adolescent human age of  $\geq 12$  years, a microscopic bone dystrophy finding, considered transient and reversible, was noted, and exposure margins at which no bone finding was noted were 5.7 to 6.1 times the human AUC at the maximum recommended human dose (MRHD) of 200 mg. No bone findings were observed in rats at any dose in the 6-month toxicity study (up to 25 times the human AUC at the MRHD of 200 mg) or in any of the toxicity studies in cynomolgus monkeys (comparable to human age of  $\geq 8$  years; up to 30 times the human AUC at the MRHD of 200 mg).

#### Genotoxicity

Abrocitinib was not mutagenic in the bacterial mutagenicity assay (*Ames* assay). It was not aneugenic or clastogenic based on the results of the *in vivo* rat bone marrow micronucleus assay.

#### Carcinogenicity

No evidence of tumorigenicity was observed in the 6-month Tg.rasH2 mice administered abrocitinib at oral doses up to 75 mg/kg/day and 60 mg/kg/day in female and male mice, respectively. In the 2-year carcinogenicity study, higher incidence of benign thymoma was noted in female rats at the lowest dose tested. Thus, a lowest observed adverse effect level (LOAEL) is set in females at exposures equal to 0.6 times the human AUC at the MRHD of 200 mg. In males the no observed adverse effect level (NOAEL) was set at exposures equal to 13 times the human AUC at the MRHD of 200 mg. The human relevance of benign thymoma is unknown.

#### Reproductive and developmental toxicity

Abrocitinib had no effects on male fertility or spermatogenesis. Abrocitinib resulted in effects on

female fertility (lower fertility index, corpora lutea, implantation sites and post-implantation loss) but no fertility effects were noted at exposures equal to 1.9 times the human AUC at the MRHD of 200 mg. The effects reversed 1 month after cessation of treatment.

No foetal malformations were observed in embryo-foetal development studies in rats or rabbits. In an embryo-foetal development study in pregnant rabbits, effects on embryo-foetal survival were noted at the lowest dose tested with exposures equal to 0.14 times the unbound human AUC at the MRHD of 200 mg. Increased litter incidences of unossified hindlimb phalanges and tarsals and forelimb phalanges were observed with effects on forelimb phalanges noted at exposures equal to 0.14 times the unbound human AUC at the MRHD of 200 mg.

In an embryo-foetal development study in pregnant rats, while increased embryo-foetal lethality was noted, none was observed at exposures equal to 10 times the human AUC at the MRHD of 200 mg. Increased incidence of skeletal variations of short 13th ribs, reduced ventral processes, thickened ribs, and unossified metatarsals were noted in the foetuses, but none were observed at exposures equal to 2.3 times the human AUC at the MRHD of 200 mg.

In a pre- and postnatal development study in pregnant rats, dams had dystocia with prolonged parturition, offspring had lower body weights and lower postnatal survival. No maternal or developmental toxicity was observed in either dams or offspring at exposures equal to 2.3 times the human AUC at the MRHD of 200 mg.

Administration of abrocitinib to juvenile rats beginning on postnatal Day 10 (comparable to a 3-month-old human infant) resulted in adverse microscopic and macroscopic bone findings, including malrotated paws, fractures, and/or femoral head abnormalities at exposures  $\geq 0.8$  times the human AUC at the MRHD of 200 mg. Administration of abrocitinib to juvenile rats beginning on postnatal Day 21 and older (comparable to a 2-year-old human and older) was not associated with microscopic or macroscopic bone findings.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1. List of Excipients**

#### **Tablet core**

LPD Title: Abrocitinib - Cibinqo

LPD rev no.: 3.1

LPD Date: August 01, 2024

Country: Thailand

Reference EU SmPC effective date: March 21, 2024

Microcrystalline cellulose (E460i)

Calcium hydrogen phosphate anhydrous (E341ii)

Sodium starch glycolate

Magnesium stearate (E470b)

### **Film-coat**

Hypromellose (E464)

Titanium dioxide (E171)

Lactose monohydrate

Macrogol (E1521)

Triacetin (E1518)

Iron red oxide (E172)

### **6.2. Incompatibilities**

Not applicable.

### **6.3. Shelf Life**

Shelf-life for CIBINQO 50 mg: 24 months

Shelf-life for CIBINQO 100 and 200 mg: 30 months

When stored at controlled room temperature for product packaged in high-density polyethylene (HDPE) bottles and polypropylene closures with heat induction seal liners and in PVDC/foil blisters.

### **6.4. Special Precautions for Storage**

This product does not have any special storage restrictions.

### **6.5. Nature and Contents of Container**

#### **CIBINQO 50, 100 and 200 mg**

High-density polyethylene (HDPE) bottle and polypropylene closure containing 14 or 30 film-

LPD Title: Abrocitinib - Cibinqo

LPD rev no.: 3.1

LPD Date: August 01, 2024

Country: Thailand

Reference EU SmPC effective date: March 21, 2024

coated tablets.

Polyvinylidene chloride (PVDC) blister with aluminum foil lidding film containing 7 film-coated tablets.

Each pack contains 7, 14 and 28 film-coated tablets.

## **7. MARKETING AUTHORIZATION HOLDER**

Pfizer (Thailand) Limited

## **8. MARKETING AUTHORIZATION NUMBERS**

CIBINQO 50 mg: 1C 15110/65 (NC)

CIBINQO 100 mg: 1C 15111/65 (NC)

CIBINQO 200 mg: 1C 15112/65 (NC)

Not all presentations or pack sizes may be marketed.

## **9. DATE OF AUTHORIZATION**

30 September 2022

## **10. DATE OF REVISION OF THE TEXT**

1 August 2024

LPD Revision No.: 3.1

LPD Date: August 01, 2024

Country: Thailand