

Prescription Only

Each soft capsule contains: Benzonatate 100mg

[Excipients]

Gelatin 160, Glycerin, Purified Water

[Pharmacology]
Benzonatate Alvogen Soft Capsules 100mg is a non-narcotic oral antitussive agent.

Benzonatate Alvogen Soft Capsules 100mg acts peripherally by

anesthetizing the stretch receptors located in the respiratory passages, lungs, and stretch receptor on pleura by dampening their activity and thereby reducing the cough reflex at its source. It begins to act within 15 to 20 minutes and its effect for 3 to 8 hours. It has no inhibitory effect on the respiratory center in recommended dosage.

#### [Indication]

For the symptomatic relief of cough.

#### [Contraindications]

Hypersensitivity to benzonatate or related compounds.

### [Warnings and Precautions]

- . Children below age 14 and those who have difficulty swallowing should not use.
- 2. Active ingredient release from the capsule in the mouth can produce a temporary local anesthesia of the oral mucosa and choking could occur. Swallow whole. Do not chew.
  Pregnancy: It should be given to a pregnant woman only if
  clearly needed. (Animal reproduction studies have not been
- conducted.) Nursing mothers: It is not known whether this drug is excreted

# in human milk. Close attention when administe Carcinogenicity, mutagenicity and reproduction studies have not been conducted.

#### [Side Effects]

Sedation, headache, slight dizziness, itching, rash, stuffy nose, constipation, nausea, gastrointestinal upset, burning eyes, fear of cold, chest pain, allergies have been reported.

#### [Overdosage]


No clear case has been reported. The drug is chemically related to tetracaine and other topical anesthetics and shares various aspects of their pharmacology and toxicology. Drugs of this type are generally well absorbed after ingestion.

[Toxic Symptoms]

If capsules are chewed or dissolved in the mouth, oropharyngeal anesthesia will develop rapidly, which may cause choking and airway compromise. Further central inhibition may lead to clonic

#### [Toxic Treatment]

Evacuate gastric contents and administer copious amounts of activated charcoal slurry. Even in the conscious patient, cough and gag reflexes may be so depressed as to necessitate




**VARIABLE DATA** 

LOT, MFD & EXP will be marked

during production as indicated.

\*LOT NUMBER format defined

by supplier so not required here



## COPYRIGHTS / CONTACT

This artwork and its content is copyright of Alvogen ©Alvogen 2012. All rights reserved.

Benzonatate Thailand 100mg x500

03

**CMYK** 

SUPPORT COLOURS (NO PRINT)

Die Cut

N/A

N/A

MINIMUM FONT SIZE = 7PT

-40 Tracking 80% horizzontal scaling

TH TEXT = 11pt

If the above minimum font size is not

acceptable for this artwork (excluding

codes) please do not approve.

THERAPEUTIC GROUP

FORM ICON

Variable data info

244x64 mm/105x64mm

DE 21/07/2020

DE 31/07/2020

ARTDB018352L

**Bottle LABEL** 

Die Cut size

Proof round **ART Code** 

PRINT COLOURS

Created Modified

Any Anthropoidea digitally or hardcopy redistribution, alternation or reproduction of part or all of the contents in any form is prohibited by other than Alvogen staff for company usage.

## Main contact:

David Garcia, Artwork Director david.garcia@Alvogen.com

