BAXTER CONFIDENTIAL - INTERNAL USE ONLY					
Part Number: BE-35	-04-112	NET	Date: 270CT20	Proofread No.: P05	
Designer: RK	DF	KHI.	Page: 1		
Colour Reference:	P185U				

AVDB3WH1G

Baxter



With **Electrolytes**

For single use only

Read the package leaflet before use Keep out of the sight and reach of children

Do not freeze

Store in the overpouch

Do not infuse before, simultaneously or after administration of blood through the same equipment

Use only if the amino acid and glucose solutions are clear, the lipid emulsion and the final mixture after reconstitution are homogenous and with a milky appearance and the container undamaged

Do not connect in series

Do not infuse without opening 1500 the seals and mixing the contents of the 3 compartments

After reconstitution, immediately administer the product

1000

ยาอันตราย

800

600

400

200

ฉลากถุงบรรจุยา ตำรับยาชื่อ OLIMEL N12E ขนาดบรรจุ 2000 ml

歐立美N12E 輸注乳液 **OLIMEL N12E**

12 g/l nitrogen 950 kcal/l with electrolytes **Emulsion for Infusion**

僅供中央靜脈輸注

For central intravenous administration

2,000 mL of the final mixture provides the following:

Total calories approx.	1,900 kcal
Non-protein calories	1,280 kcal
Nitrogen	24.0 g
Non-protein calories /	
nitrogen ratio	53 kcal/g
Glucose	146.7 g
Lipids	70.0 g
Glucose / lipid calories ratio	45/55
Amino acids	151.9 g
Glucose calories	580 kcal
Lipid calories	700 kcal
Lipid / total calories	37%
Electrolytes:	
Sodium	70.0 mmol
Potassium	60.0 mmol
Magnesium	8.0 mmol
Calcium	7.0 mmol
Phosphate	29.2 mmol
Acetate	140 mmol
Chloride	90 mmol
рН	6.4
Osmolarity	1,270 m0sm/L

ขอรับรองว่าข้อความบนฉลากของ ภาชนะบรรจุทุกชนิดและหีบห่อ ถกต้องตรงตามมาตรา ๒๕ (๓) หรือ ๒๗ (๒) และ (๓) แล้วแต่กรณี

2,000 mL

Nitrogen g

Total kcal

Non-protein calories

24.0

1.900

1,280

Baq

Formula for 2,000 mL after mixing:

Refined olive oil + refined soybean oil 70.00 g, Alanine 21.97 g, Arginine 14.88 g. Aspartic acid 4.39 g. Glutamic acid 7.58 g, Glycine 10.53 g, Histidine 9.06 g, Isoleucine 7.58 g, Leucine 10.53 g, Lysine (equivalent to lysine acetate) 11.95 g (16.85 g), Methionine 7.58 g, Phenylalanine 10.53 g, Proline 9.06 g, Serine 5.99 g, Threonine 7.58 g, Tryptophan 2.53 g, Tyrosine 0.39 g, Valine 9.72 g, Sodium acetate, trihydrate 2.99 g, Sodium glycerophosphate, hydrated 7.34g, Potassium chloride 4.47 g, Magnesium chloride, hexahvdrate 1.62 g. Calcium chloride, dihydrate 1.03 g, Glucose (equivalent to glucose monohydrate) 146.67 g (161.33 g)

Excipients:


purified egg phosphatide, glycerol, sodium oleate, sodium hydroxide, water for injections, glacial acetic acid, hydrochloric

After reconstitution: Emulsion for Infusion 2,000 mL in a 3-compartment bag Glucose solution 27.5%: 533 mL Amino acid solution 14.2%: 1,067 mL Lipid emulsion 17.5%: 400 mL

百特醫療產品股份有限公司 台北市敦化南路二段95號28樓 製造廠:Baxter S.A 廠址:Boulevard René Branquart, 80, 7860 Lessines, Belgium 適應症、用法用量及注意事項: 請詳見仿單 請勿冷凍

Manufactured by: Baxter SA, 7860 Lessines, Belgium Imported by: Baxter Healthcare (Thailand) Co, Ltd., Bangkok, Thailand Thailand Reg. No. 2C xx/6x







AVDB3WH1G

ยาสินอายุ



BE-35-04-112

Mfg. Date:

Lot: