

ฉลากซองบรรจุยา
(ขนาดบรรจุ 3.5 กรัม)

ด้านหน้า



The Original **Calmoseptine[®]** Ointment

Reg. No. : 2C XX/XX Net wt. 1/8oz (3.5g)

ยาใช้ภายนอก
ห้ามรับประทาน

MFG DD/MM/YY
EXP DD/MM/YY

7 46876 00000 2
Lot. no.

MADE IN USA Manufactured by: Calmo Manufacturing Inc., Huntington Beach, CA 92647
Importer: Zuellig Pharma Ltd. Bangkok, Thailand **Store between: 15-30°C (59-86°F)**

Anorectal Ointment • temporarily protects irritated areas
• temporary relief of pain or burning and skin irritations

ด้านหลัง

Drug Facts		Drug Facts (continued)
Active ingredients	Purpose	<ul style="list-style-type: none">• Do not put this product into the rectum by using fingers or any mechanical device or applicator.• Do not exceed the recommended daily dosage unless directed by a doctor.
Menthol 0.44%.....Anorectal Analgesic Zinc Oxide 20.6%....Anorectal Protectant, Astringent		Stop use and ask a doctor if <ul style="list-style-type: none">• condition worsens or does not improve within 7 days • bleeding occurs • symptom being treated does not subside or if redness, irritation, swelling, pain or other symptoms develop or increase. Keep out of reach of children. In case of accidental ingestion contact a Poison Control Center immediately.
Uses • Temporarily protects irritated areas. • For the temporary relief of: • local itching and discomfort in the perianal area • pain or burning • skin irritations		Directions Adults: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth or air dry before application of this product. • Apply externally to the affected area up to 6 times daily or after each bowel movement. • Children under 12 years of age: consult a doctor.
Warnings: For external use only		Inactive ingredients chlorothymol, glycerin, lanolin, petrolatum, phenol, sodium bicarbonate, and thymol in a suitable ointment base
When using this product <ul style="list-style-type: none">• Certain persons can develop allergic reactions to ingredients in this product.• Avoid contact with eyes.		

Thai Anorectal Sachet Artwork 11-30-2020v1 Outlined

ขอรับรองว่าข้อความบนฉลากของภาชนะบรรจุ
ทุกชนิดและหีบห่อถูกต้องตรงตามมาตรา ๒๕ (๓)
หรือ ๒๗ (๒) และ (๓) แล้วแต่กรณี

38mm

78mm

38mm

78mm

ฉลากกล่องยา
(ขนาดบรรจุ 20 ซอง)

Reg. No. : 2C XX/XX



The Original

NDC 0799-0001-05

Net wt. 1/8oz (3.5g)

Calmoseptine[®] Ointment

Anorectal Ointment • temporarily protects irritated areas
• temporary relief of pain or burning and skin irritations

20 count - 3.5 gram packets

Store between: 15-30°C (59-86°F)

FOR EXTERNAL USE ONLY

ยาใช้ภายนอก
ห้ามรับประทาน

MADE IN USA

Manufactured by: Calmo Manufacturing, Inc.
16601 Burke Lane
Huntington Beach, CA 92647

1-800-800-3405
www.calmoseptine.com

Imported by:
Zuellig Pharma Ltd.
Bangkok, Thailand

Lot. no.
MFG DATE: DD/MM/YY
EXP DATE: DD/MM/YY
ยาสิ้นอายุ

ขอรับรองว่าข้อความบนฉลากของภาชนะบรรจุ
ทุกชนิดและหีบห่อถูกต้องตรงตามมาตรา ๒๕ (๓)
หรือ ๒๗ (๒) และ (๓) แล้วแต่กรณี

Active ingredients

Menthol 0.44%.....Anorectal Analgesic
Zinc Oxide 20.6%.....Anorectal Protectant, Astringent

Purpose

Uses: • Temporarily protects irritated areas.
• For the temporary relief of: • local itching and discomfort in the perianal area • pain and burning • skin irritations

Warnings: For external use only

When using this product

• Certain persons can develop allergic reactions to ingredients in this product. • Do not put this product into the rectum by using fingers or any mechanical device or applicator. • Do not exceed the recommended daily dosage unless directed by a doctor. • Avoid contact with eyes.

Stop use and ask a doctor if

• condition worsens or does not improve within 7 days • bleeding occurs
• symptom being treated does not subside or if redness, irritation, swelling, pain or other symptoms develop or increase.

Keep out of reach of children. In case of accidental ingestion contact a Poison Control Center immediately.

Directions: Adults: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth or air dry before application of this product. • Apply externally to the affected area up to 6 times daily or after each bowel movement. • Children under 12 years of age: consult a doctor.

Inactive Ingredients chlorothymol, glycerin, lanolin, petrolatum, phenol, sodium bicarbonate, and thymol in a suitable ointment base

38mm

95mm

38mm