

BAXTER CONFIDENTIAL - INTERNAL USE ONLY		
Part Number: BE-35-04-111	Date: 27OCT20	Proofread No.: P05
Designer: RK	Page: 1	
Colour Reference:	P185U	

AVDB3WH1F

Baxter

Central Vein
Only

With
Electrolytes

For single use only

Read the package leaflet before use

Keep out of the sight and reach of children

Do not freeze

Store in the overpouch

Do not infuse before, simultaneously or after administration of blood through the same equipment

Use only if the amino acid and glucose solutions are clear, the lipid emulsion and the final mixture after reconstitution are homogenous and with a milky appearance and the container undamaged

Do not connect in series

Do not infuse without opening the seals and mixing the contents of the 3 compartments

After reconstitution, immediately administer the product

1,000

ยาอันตราย

500

ฉลากถุงบรรจุยา ตำรับยาชื่อ OLIMEL N12E

ขนาดบรรจุ 1500 ml

“百特” 歐立美N12E 輸注乳液
OLIMEL N12E

12 g/l nitrogen 950 kcal/l
with electrolytes,
Emulsion for Infusion

僅供中央靜脈輸注

For central intravenous administration

1,500 mL of the final mixture provides the following:

Total calories approx.	1,420 kcal
Non-protein calories	960 kcal
Nitrogen	18.0 g
Non-protein calories / nitrogen ratio	53 kcal/g
Glucose	110.0 g
Lipids	52.5 g
Glucose / lipid calories ratio	45/55
Amino acids	113.9 g
Glucose calories	430 kcal
Lipid calories	520 kcal
Lipid / total calories	37%
Electrolytes:	
Sodium	52.5 mmol
Potassium	45.0 mmol
Magnesium	6.0 mmol
Calcium	5.3 mmol
Phosphate	21.9 mmol
Acetate	105 mmol
Chloride	68 mmol
pH	6.4
Osmolarity	1,270 mOsm/L

AVDB3WH1F

(01)00085412676869



ยาสิ้นอายุ
Exp:

ขอรับรองว่าข้อความบนฉลากของ
ภาชนะบรรจุทุกชนิดและหีบห่อ
ถูกต้องตรงตามมาตรา ๒๕ (๓)
หรือ ๒๗ (๒) และ (๓) แล้วแต่กรณี

John S. Kim

1,500 mL

Bag

Nitrogen g	18.0
Total kcal	1,420
Non-protein calories	960

Formula for 1,500 mL after mixing:

Refined olive oil + refined soybean oil 52.50 g, Alanine 16.48 g, Arginine 11.16 g, Aspartic acid 3.30 g, Glutamic acid 5.69 g, Glycine 7.90 g, Histidine 6.79 g, Isoleucine 5.69 g, Leucine 7.90 g, Lysine (equivalent to lysine acetate) 8.96 g (12.64 g), Methionine 5.69 g, Phenylalanine 7.90 g, Proline 6.79 g, Serine 4.50 g, Threonine 5.69 g, Tryptophan 1.90 g, Tyrosine 0.30 g, Valine 7.29 g, Sodium acetate, trihydrate 2.24 g, Sodium glycerophosphate, hydrated 5.51 g, Potassium chloride 3.35 g, Magnesium chloride, hexahydrate 1.22 g, Calcium chloride, dihydrate 0.77 g, Glucose (equivalent to glucose monohydrate) 110.00 g (121.00 g)

Excipients:

purified egg phosphatide, glycerol, sodium oleate, sodium hydroxide, water for injections, glacial acetic acid, hydrochloric acid.

After reconstitution: Emulsion for Infusion 1,500 mL in a 3-compartment bag
Glucose solution 27.5% : 400 mL
Amino acid solution 14.2% : 800 mL
Lipid emulsion 17.5% : 300 mL

百特醫療產品股份有限公司
台北市敦化南路二段95號28樓
製造廠:Baxter S.A
廠址:Boulevard René Branquart, 80, 7860
Lessines, Belgium


適應症、用法用量及注意事項:
請詳見仿單
請勿冷凍

Manufactured by: Baxter SA, 7860
Lessines, Belgium
Imported by: Baxter Healthcare
(Thailand) Co, Ltd.,
Bangkok, Thailand
Thailand Reg. No. 2C xx/6x

POM

BE-35-04-111

Lot:



Mfg. Date: