





## 60 doses

## Salflumix Easyhaler®

50 micrograms/ 250 micrograms/dose, inhalation powder

salmeterol 50 microg /fluticasone propionate 250 microg Inhalation use.

Read the package leaflet before use.

Use within 1 month of removing from the foil bag.



MFG:

MFG DATE

EXP:

**EXPDATE** 

LOT:

000000-0

ขอรับรองข้อความบนฉลากของภาชนะบรรจุทุกชนิดและหี้บห่อ ถูกต้องตามมาตรา ๒๕ (๓) หรือ ๒๗ (๒) และ ๒๗ (๓) สี่ของฉลากพื้นสีขาว อักษรสีดำ


18.07.2022/KS


Min. font 7 pt

**PROOFBOX** 

size 48x80 mm

ซึ่งอยู่บนแผ่นเดียวกัน





## ฉลากขวดยา (เหมือนกันทุกขนาดบรรจุ)



64,3

ขอรับรองข้อความบนฉลากของภาชนะบรรจุทุกชนิดและหืบห่อ ถูกต้องตามมาตรา ๒๕ (๓) หรือ ๒๗ (๒) และ ๒๗ (๓)

สีของฉลากพื้นสีขาว เขียว อักษรสีดำ ซึ่งอยู่บนแผ่นเดียวกัน

27



DATE	14.07.2022	SB-REVISION	01	VERSION	1.0
CLIENT	Orionpharma			FORMAT	HD 16:9
ID				LANGUAGE	English (Master)
subject จาก QR code บนกล่อง				deo แนะนำวิธีการใช้ยา ตาม	

## Salflumix Easyhaler® APAC English Story board 1

หมายเหตุ - QR code ยังไม่ Valid จนกว่าทะเบียนจะได้รับการอนุมัติ

[Storyboard]

	On-screen text	Pictures	Speaker's Text
L	This video is intended for patients who have been prescribed Salflumix Easyhaler®.  1.0	TEASYHALER®  This video is intended for patients who have been prescribed Salflumix Easyhaler®.	
<u> </u>		Sallumik EASYHALER	This device is a metered dose, dry powder inhaler for the treatment of respiratory diseases like asthma or COPD. The inhaler comes in a foil bag. Take the inhaler out of it only when you are ready to start using it for the first time.

	On-screen text	Pictures	Speaker's Text
03			There is a dust cap for the inhaler which prevents the inhaler from preloading.
04		Estation	Before you start taking a dose remove the dust cap. Thereafter, place the inhaler between your forefinger and your thumb.
05	Shake	Shake	Shake the inhaler in an upright position 3-5 times.

	On-screen text	Pictures	Speaker's Text
	Click	Click	Press down until you hear a click. Then let it go back to its original position.
,	Inhale	Inhale	Breathe out normally, place the mouthpiece in your mouth, between your teeth and close your lips tightly around the mouthpiece. Take a strong deep breath. Take the inhaler out of your mouth, hold your breath for at least 5 seconds. Then breathe out normally. If you need to take more than one dose, repeat the steps from Shake onwards.  After all doses put the dust cap back on the mouthpiece.
•	Rinse Spit it out		After taking your dose, rinse your mouth with water and spit it out.
		Rinse	