
DOCUMENT INFORMATION	
Document Name	FJ_Ef_TH_10_IDT_Version_05
Date	17.10.2017
Article No.	

ARTICLE INFORMATION	
Product	Ferinject
Manufacturer	IDT
Country	Thailand
Language(s)	English, Thai
Presentation Form	Vials
Content	10 ml
Strength	50 mg/ml

DESIGN AND FONTS	
Fonts	Arial, Mic32
Font Sizes	min: 5 pt max: 6.7 pt
Logos	Ferinject, Vifor Pharma
Price	
Other Labelling Requirements	

Colours/Flats	
Black Pantone 485 C	
Pantone 429 U	

MANUFACTURER PACKAGING SPECIFICATIONS	
Manufacturer Machine Code and Type	
Manufacturer Identification Number	
Information in 2D Code	
Format and Content Imprints Variable Data	Lot = Lot: Man. Date = Man.date: Exp = Expiry/ยาสิ้นอายู:
Dimensions	26 x 73 mm
Onesided or double-sided printing	double




DOCUMENT INFORMATION	
Document Name	FJ_Kf_TH_1x10_IDT_V10
Date	30.10.2018
Article No.	

ARTICLE INFORMATION	
Product	Ferinject
Manufacturer	IDT
Country	Thailand
Language(s)	English, Thai
Presentation Form	Vials
Content	1x10 ml
Strength	50 mg/ml

DESIGN AND FONTS	
Fonts	Arial
Font Sizes	min: 5.4 pt max: 13 pt
Logos	Ferinject, Vifor Pharma
Product Code (PC)	XXXXXXXXXX
Braille	n/a
Price	n/a
Registration Number	
Other Labelling Requirements	EAN13 based on PC

Colours/Flats	
Black	Pantone 485 C
Pantone 254 C	Lacquer Free
Dye cut	

MANUFACTURER PACKAGING SPECIFICATIONS	
Manufacturer Machine Code and Type	
Manufacturer Identification Number	
Information in 2D Code	
Tamper evidence	Safety Seal
Format and Content Imprints Variable Data	Lot = Lot: Man. Date = Man.date: Exp = Expiry/ยาสิ้นอายุ:
Dimensions	37x37x78 mm





ferric carboxymaltose

50 mg iron/ml

solution for injection/infusion

Imported by: Zuellig Pharma Ltd. 8–9/F Ploenchit Center, 2 Sukhumvit Rd., Kwaeng Klongtoey, Khet Klongtoey, Bangkok 10110

Manufactured and packed by: IDT Biologika GmbH. Dessau Rosslau/Germany

Released by: Vifor (International) Inc. St.Gallen/Switzerland



ferinject[®]

ferric carboxymaltose

Before using check on absence of sediments and observe

expiry date.
For administration by IV
infusion the solution should be diluted in sterile 0.9% m/V sodium chloride solution. Use immediately after dilution.

Each mI contains 50 mg iron (as ferric carboxymaltose). Water for injections, sodium hydroxide, hydrochloric acid. For intravenous use. Read the package leaflet before

Kead the package leaflet before use.

Keep out of the reach and sight of children.

Store in the original carton.

Do not store above 30 °C.

Do not freeze.

500 mg

Kf Thailand 1x10 ml IDT-Materialnummer 37/37/78





ferinject[®]

ferric carboxymaltose 50 mg iron/ml solution for injection/infusion

1 vial (10 ml)

One 10 ml vial



Reg. No.:



ยาควบคุมพิเศษ ใช้เฉพาะสถานพยาบาล



contains 500 mg iron

500 mg

500 mg







