

BAXTER CONFIDENTIAL - INTERNAL USE ONLY		
Part Number: BE-35-04-109	Date: 27OCT20	Proofread No.: P05
Designer: RK	Page: 1	
Colour Reference:	P185U	

AVDB3WH1C
Baxter

Central Vein
Only

With
Electrolytes

For single use only

Read the package leaflet before use
Keep out of the sight and reach of
children

Do not freeze

Store in the overpouch

Do not infuse before,
simultaneously or after administration
of blood through the same equipment

Use only if the amino acid and
glucose solutions are clear,
the lipid emulsion and the final
mixture after reconstitution are
homogenous and with a milky
appearance and the container
undamaged

Do not connect in series

Do not infuse without opening
the seals and mixing the contents of
the 3 compartments

After reconstitution,
immediately administer the
product

300



ยาอันตราย

Mfg. Date:

ฉลากถุงบรรจุยา ตำรับยาชื่อ OLIMEL N12E
ขนาดบรรจุ 650 ml

“百特” 歐立美N12E 輸注乳液
OLIMEL N12E

12 g/l nitrogen 950 kcal/l with electrolytes
Emulsion for Infusion

For central intravenous administration 僅供中央靜脈輸注
650 mL of the final mixture provides the following:

Total calories approx.	620 kcal
Non-protein calories	420 kcal
Nitrogen	7.8 g
Non-protein calories / nitrogen ratio	53 kcal/g
Glucose	47.7 g
Lipids	22.8 g
Glucose / lipid calories ratio	45/55

Amino acids	49.4 g	Electrolytes:	
Glucose calories	190 kcal	Sodium	22.8 mmol
Lipid calories	230 kcal	Potassium	19.5 mmol
Lipid / total calories	37%	Magnesium	2.6 mmol
		Calcium	2.3 mmol
		Phosphate	9.5 mmol
		Acetate	46 mmol
		Chloride	30 mmol
		pH	6.4
		Osmolarity	1,270 mOsm/L

AVDB3WH1C

(01)00085412676296



ยาอันตราย
Exp:

ขอรับรองว่าข้อความบนฉลากของ
ภาชนะบรรจุทุกชนิดและหีบห่อ
ถูกต้องตรงตามมาตรา ๒๕ (๓)
หรือ ๒๗ (๒) และ (๓) แล้วแต่กรณี

John S.M.

650 mL

Nitrogen g
Total kcal

Per bag

7.8
620
420

Non-protein calories

Formula for 650 mL after mixing:
Refined olive oil + refined soybean oil
22.75 g, Alanine 7.14 g, Arginine
4.84 g, Aspartic acid 1.43 g, Glutamic acid
2.46 g, Glycine 3.42 g, Histidine 2.94 g,
Isoleucine 2.46 g, Leucine 3.42 g, Lysine
(equivalent to lysine acetate) 3.88 g
(5.48 g), Methionine 2.46 g, Phenylalanine
3.42 g, Proline 2.94 g, Serine 1.95 g,
Threonine 2.46 g, Tryptophan 0.82 g,
Tyrosine 0.13 g, Valine 3.16 g, Sodium
acetate, trihydrate 0.97 g, Sodium
glycerophosphate, hydrated 2.39 g,
Potassium chloride 1.45 g, Magnesium
chloride, hexahydrate 0.53 g, Calcium
chloride, dihydrate 0.34 g, Glucose
(equivalent to glucose monohydrate)
47.67 g (52.43 g)

Excipients: purified egg phosphatide,
glycerol, sodium oleate, sodium hydroxide,
water for injections, glacial acetic acid,
hydrochloric acid.

After reconstitution: Emulsion for Infusion

650 mL in a 3-compartment bag
Glucose solution 27.5% : 173 mL

Amino acid solution 14.2% : 347 mL

Lipid emulsion 17.5% : 130 mL

百特醫療產品股份有限公司

台北市敦化南路二段95號28樓

製造廠: Baxter S.A

廠址: Boulevard René Branquart, 80, 7860

Lessines, Belgium

適應症、用法用量及注意事項:

請詳見仿單

請勿冷凍

Manufactured by: Baxter SA, 7860

Lessines, Belgium

Imported by: Baxter Healthcare

(Thailand) Co, Ltd.,

Bangkok, Thailand

Thailand Reg. No. 2C xx/6x

BE-35-04-109

Lot:

POM

